

# Mapping the distribution of armored harvestmen (Opiliones, Laniatores) in Colombia: updated list of species, taxonomic contributions, and insight of diversity in protected areas

Osvaldo Villarreal<sup>1,2</sup>, Daniela Ahumada-C.<sup>3,4</sup>, Leonardo Delgado-Santa<sup>5</sup>

<sup>1</sup> Departamento de Invertebrados, Museu Nacional/Universidade Federal do Rio de Janeiro, Quinta da Boa Vista, São Cristóvão, 20.940-040, Rio de Janeiro, RJ, Brazil

<sup>2</sup> Museo del Instituto de Zoología Agrícola, Facultad de Agronomía, Universidad Central de Venezuela, Maracay, Aragua, Venezuela

<sup>3</sup> Grupo de Investigación Biología Descriptiva y Aplicada, Programa de Biología, Universidad de Cartagena, Cartagena de Indias, Bolívar, Colombia

<sup>4</sup> Grupo de Investigación Hidrobiología, Programa de Biología, Universidad de Cartagena, Cartagena de Indias, Bolívar, Colombia

<sup>5</sup> Grupo de Investigación Ecdysis, Programa de Biología, Universidad del Quindío, Armenia, Quindío, Colombia

Corresponding authors: Leonardo Delgado-Santa ([leonardodelgadosanta@gmail.com](mailto:leonardodelgadosanta@gmail.com)); Osvaldo Villarreal ([osvaldovillarreal@gmail.com](mailto:osvaldovillarreal@gmail.com))



Academic editor: Adriano Kury

Received: 22 February 2023

Accepted: 11 July 2023

Published: 17 August 2023

ZooBank: <https://zoobank.org/A5C1D3DF-FF87-404A-9EA0-4608CF89EFA7>

Citation: Villarreal O, Ahumada-C. D, Delgado-Santa L (2023) Mapping the distribution of armored harvestmen (Opiliones, Laniatores) in Colombia: updated list of species, taxonomic contributions, and insight of diversity in protected areas. ZooKeys 1175: 223–284. <https://doi.org/10.3897/zookeys.1175.102485>

Copyright: © Osvaldo Villarreal et al.  
This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International – CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

## Abstract

Colombia is a biodiverse country with 1,486 protected areas covering almost 50 million hectares, yet little is known about the biodiversity they harbor, particularly in terms of harvestmen (Arachnida: Opiliones). This study provides a comprehensive updated summary of the armored harvestmen (Laniatores) species found in Colombia with a list of 173 nominal species, focusing on the Laniatores fauna found in protected areas and its diversity is detailed and analyzed. Maps with all records associated with ecoregions and protected areas in Colombia are presented. Additionally, three new Laniatores species are described here: *Phalangodus andresi* **sp. nov.** from the department of Cundinamarca, and *Ventrifurca phallaina* **sp. nov.** and *Ampycella fortunata* **sp. nov.** from the department of Valle del Cauca; and a new family assignment is proposed for *Paraphalangodus* Roewer, 1915, placing it in the family Nomoclastidae. Information available on Laniatores in the National System of Protected Areas is still scarce and promoting strategies to facilitate the regulatory procedures for collecting specimens in these areas and increasing investment in basic science projects, are suggested to improve the understanding and study of the Laniatores fauna and other invertebrates in Colombia. Finally, a chronicle and timeline set of figures of species of Laniatores from Colombia, described by various authors during three periods, is given.

**Key words:** Checklist, ecoregions, harvestmen, Neotropics, protected natural areas

## Introduction

Colombia is one of the most biodiverse countries in the world (Andrade-C 2011). This diversity is represented and legally protected in 1,486 conservation units (covering 49,884,326.77 hectares), under the figure of National System of Protected Areas (RUNAP 2022). These areas allow ecological self-regulation, their ecosystems have not been substantially altered by human exploitation or occupation, and their biodiversity is under management (CONPES 2021). Although Colombia is one of the countries in South America with the largest



number of areas under conservation (CONPES 2021), knowledge about the species it harbors is scarce, especially in terms of harvestmen (Arachnida: Opiliones). Colombia ranks third in South America in terms of the number of recorded Opiliones species, only behind Brazil (1,010) and Venezuela (392) (Villarreal et al. 2021a). The first effort to list the arachnid species of Colombia was made by Flórez and Sánchez-C (1995), summarizing 77 species of harvestmen. Later, all records and distribution data on Colombian Laniatores, which constitute the vast majority of Opiliones in the country, were compiled by Kury (2003) into an excellent New World Laniatores catalog. Taxonomic studies on harvestmen have become increasingly frequent in the Neotropics, especially in Colombia, where numerous genera and species have been recently described or recorded (e.g., Villarreal and Rodríguez 2006; García 2014; Villarreal and García 2016; Pinzón and Pinto-da-Rocha 2020; Villarreal and García 2021). More recently, a general diagnosis of the current knowledge of the group in Colombia was published (Perafán et al. 2013), recording a total of 162 Opiliones species in the country. Minor local inventory works mention 170 (García and Medrano 2015) and 186 (Barriga et al. 2019) species of Opiliones in Colombia. Only two opilionofaunal inventory and/or ecological studies have been conducted in Colombia to our knowledge so far (García and Medrano 2015; De Moya et al. 2021). García and Medrano (2015) carried out the guide of Opiliones of the Reserva Natural Rio Nambí, a protected area on the coastal foothills (department of Nariño) (CONPES 2021), recording 12 families and a wealth of undescribed species. More recently, De Moya et al. (2021) studied the diversity of Opiliones by altitude in Sierra Nevada de Santa Marta (department of Magdalena), an area of high conservation importance due to its significant climatic variability and high rates of endemism, recording nine families. Studies focused on analyzing the composition of neotropical harvestmen in protected areas are not frequent. These have been carried out mainly in Brazil, among them, Soares and Soares (1970) describe seven species for the Itatiaia Biological Station (Rio de Janeiro); Adis et al. (2002) record seven species for the Ducke Reserve (Amazonia), and Bragagnolo and Pinto-da-Rocha (2003) recorded 52 species in the Serra dos Órgãos National Park (Rio de Janeiro). A recent precedent was made by Guerrero (2019), who recorded 17 species from 30 protected areas in the province of Buenos Aires (Argentina). However, like the situation in Colombia, he concludes that many important areas still remain for the conservation of diversity that have yet to be surveyed to understand their harvestmen composition.

Since the publication of the New World Laniatores catalog by Kury (2003), more than 20 years ago, Colombia has undergone numerous changes and additions to its opilionofauna. This is partly due to the appearance of a growing group of enthusiastic local and regional arachnology researchers who have turned their interest to the fauna of this country, as well as the expansion of its biological collections and collaboration with other renowned institutions. Currently, Colombia has 162 species of armored harvestmen (Opiliones: Laniatores) recorded, belonging to 12 families of which 46 species have been recorded or described after 2003. Therefore, we consider it opportune to provide an updated and detailed summary of the taxonomic changes in the Colombian laniatorean fauna after 2003.

The purpose of this paper was to provide an updated list of species and summarize the taxonomic changes and additions that occurred after 2003. Additionally, we aim to relate the distribution of Colombian species with the biogeographical



areas proposed by WWF, discuss the presence of species in the country's National System of Protected Areas (NSPA), and present the descriptions of two new cranaid species from the departments of Cundinamarca and Valle del Cauca as well as one new gonyleptid species from the department of Valle del Cauca.

### **Taxonomic background for *Phalangodus*, *Ventrifurca*, and *Ampycella* of Colombia**

The neotropical family Cranidae has attracted attention from taxonomists in recent years; nevertheless, the definition of generic or suprageneric groups in this family remains unsatisfactory (Kury and Villarreal 2015; Villarreal and García 2016). Although some groups have been reviewed, leading to the resolution of generic synonyms, new combinations or redefinitions, improper placements, as well as descriptions of new species (e.g., Orrico and Kury 2009; Pinto-da-Rocha and Bonaldo 2011; Hara et al. 2014; Villarreal et al. 2015; Villarreal and García 2016; Hara et al. 2017), much work remains to be done.

Only three works have analyzed the phylogenetic relationships within the family Cranidae and the genus *Phalangodus* Gervais, 1842 has only been studied in a phylogenetic context in an analysis of the familial relationships of Gonyleptoidea (Kury and Villarreal 2015), where it appears as sister group to the other four representatives of the family. Despite this, the taxonomy of *Phalangodus* is relatively well understood, with a total of six species described, five of them known from Colombia (Villarreal and García 2016). A similar case exists for the genus *Ventrifurca* Roewer, 1913, which was recently reviewed (Villarreal et al. 2015), revealing some synonyms and the discovery of a new species from the department of Quindío. Currently, three species of *Ventrifurca* are known from Colombia, distributed in the western Andes.

Two species have been recorded from Colombia, belonging to the family Ampycidae, *Ampycus telifer* (Butler, 1873), and *Licornus tama* Villarreal & Kury, 2012 (García 2014). This family lacks taxonomic revisionary work, and its genera are poorly delimited or defined. Recent works have contributed to better understanding the diversity of some genera within Ampycidae (e.g., Tourinho and Mendes 2014; Hara et al. 2017) or described new species (Villarreal and Kury 2012). The genus *Ampycella* Roewer, 1929 includes two Andean species from Ecuador (Kury et al. 2022), vaguely described and illustrated and whose genital morphology is unknown.

This paper describes three species: two new cranaid (*Phalangodus* and *Ventrifurca*) and one ampycid (*Ampycella*, including the first image of the male genitalia for the genus), collected in the departments of Cundinamarca in the Central Mountain Range in the Colombian Andes and the department of Valle del Cauca in Chocó Biogeographic region.

## **Materials and methods**

### **Taxonomy**

Individuals of each species were photographed using a Leica M205C stereoscope attached to a Leica DFC450 digital camera, a Wild Heerbrugg stereoscope attached to a Nikon COOLPIX P900 and a Wild Heerbrugg microscope



attached to a HAYEAR 2307 digital camera. The multiple resultant images at different focal planes were combined with Combine ZP Suite software (Hadley 2015) to increase the depth of field and were thereafter edited in Photoshop CC 2014 software. Drawings of the species were made using Illustrator CC 2017 and Inkscape 1.2.2 softwares (Harrington 2004–2005). To color descriptions the standard names of the 267 Color Centroids of the NBS/IBCC Color System were used as named in Centore (2016). Male genitalia were studied using standard methods for this structure (Acosta et al. 2007).

Morphological terminology and patterns of taxonomic description follow Villarreal and García (2016) with slight modifications; integumentary ornamentation follows DaSilva and Gnaspini (2010); terminology for chaetotaxy of penis ventral plate follows Kury and Villarreal (2015) with the modifications proposed by Villarreal and García (2016) for the genus *Phalangodus*; the ovipositor morphology follows Villarreal and García (2016); terminology of dorsal scutum outline types follows Kury and Medrano (2016).

The first-order administrative divisions of Colombia (departments) are underlined. Maps were made using ArcGIS® 10.1 software (ESRI 2022). Colored areas represent WWF Terrestrial Eco-regions of the World (Olson et al. 2001), here abbreviated as WWF.

Morphometric abbreviations are: **AL** = maximum abdominal scutum length; **AW** = maximum abdominal scutum width; **BaCh** = basichelicerite length; **Cl** = claw; **CL** = carapace length; **ClPp** = pedipalp claw; **CW** = maximum carapace width; **DS** = dorsal scutum; **DSL** = dorsal scutum length; **Fe** = femur; **FeL I** = femur length I; **FeL II** = femur length II; **FeL III** = femur length III; **FeL IV** = femur length IV; **IOD** = inter ocular distance; **MS** = macrosetae of penis; **Mt** = Metatarsus; ; **Pp** = pedipalps; **FePp** = pedipalpal femur; **PaPp** = pedipalpal patella; **TaPp** = pedipalpal tarsus; **TiPp** = pedipalpal tibia; **Ta** = tarsus; **Ti** = tibia; **TiL I** = tibia length I; **TiL II** = tibia length II; **TiL III** = tibia length III; **TiL IV** = tibia length IV; **VP** = ventral plate. All measurements are in mm unless otherwise noted. The material studied is deposited in the arachnological collections of Instituto de Ciencias Naturales (**ICN**), Universidad Nacional, Bogotá, Colombia; Colección de Insectos de la Universidad del Quindío (**CIUQ**), Armenia, Colombia; Museo del Instituto de Zoología Agrícola “Francisco Fernández Yépez” (**MIZA**), Maracay, Venezuela; and Museu Nacional, Universidade Federal do Rio de Janeiro (**MNRJ**), Rio de Janeiro, Brazil.

## Species inventory

A comprehensive list of all described Laniatores species occurring in Colombia was compiled, based on the bibliography published until April 2023, including all valid species of Laniatores recorded for Colombia. The family and superfamily classification follows Kury et al. (2022). The logonymic information is updated only for Colombian species that were described, recorded, or underwent nomenclatural changes after New World Laniatores catalog (Kury 2003). The taxonomic references in the logonymy of the listed species were not considered in the References section.

A list of all protected areas under the NSPA jurisdiction, which harbor records of Laniatores species, was compiled. Only areas with some form of public regulation for their protection were considered. For each protected area, the families and species of armored harvestmen recorded were listed (Table 1). The source of the species data can be seen in the Suppl. material 1.



Table 1. Opiliones species by family recorded in RUNAP from Colombia.

Category	RUNAP	Family	Species
Regional Integrated Management Districts (RDIM)	Cuenca alta del Río Atrato	Manaosbiidae	<i>Camelianus fuhrmanni</i>
	de la cuenca alta del Río Quindío de Salento	Cranaidae	<i>Allocranaus columbianus</i>
		Nomoclastidae	<i>Quindina bella</i>
		Stygnidae	<i>Eutimesius ehippiatus</i>
	Delta del Río Ranchería	Cranaidae	<i>Cranaus albipustulatus</i>
	Paramo de Guargua y laguna Verde	Cosmetidae	<i>Rhaucus serripes</i>
	Paramo de guerrero	Cosmetidae	<i>Rhaucus quinquelineatus</i>
	Páramos de Guantiva y la Rusia	Cosmetidae	<i>Rhaucus papilionaceus</i>
	Río Rubachoque y pantano de arce	Cosmetidae	<i>Rhaucus vulneratus</i>
	San Miguel	Cranaidae	<i>Megacranaus pygoplus</i>
	Sector Salto del Tequendama y Cerro Manjui	Agoristenidae	<i>Muscopilio onod</i>
		Cosmetidae	<i>Eulibitia maculata</i>
			<i>Rhaucus serripes</i>
	Serranía de los Yariguies	Agoristenidae	<i>Leptostygnus yarigui</i>
		Cosmetidae	<i>Rhaucus papilionaceus</i>
		Cranaidae	<i>Phalangodus briareos</i>
	Serrania de Perija	Agoristenidae	<i>Avima troglobia</i>
			<i>Avima venezuelica</i>
National Natural Parks (NNP)	Chingaza	Cosmetidae	<i>Libitia bipunctata</i>
	Paramillo	Agoristenidae	<i>Avima tuttifrutti</i>
	Sierra de la macarena	Cosmetidae	<i>Meterginus prosopis</i>
		Cranaidae	<i>Phareicranaus angelicus</i>
	Sumapaz	Cosmetidae	<i>Eulibitia maculata</i>
			<i>Rhaucus vulneratus</i>
		Fissiphallidae	<i>Fissiphallius spinulatus</i>
Regional Natural Park (RNP)	Yaigoje Apaporis	Cosmetidae	<i>Sibambea cincta</i>
	Los Besotes	Cosmetidae	<i>Eucynorta quadripustulata</i>
	Serrania del Perija	Cosmetidae	<i>Eulibitia victoriae</i>
	Sisavita	Cosmetidae	<i>Eulibitia helena</i>
Civil Society Nature Reserve (CSNR)	La Palmita	Cosmetidae	<i>Eulibitia chacuamarei</i>
National Protective Forest Reserve (NPFR)	Bosque Oriental de Bogotá	Cosmetidae	<i>Libitia bipunctata</i>
			<i>Libitia cordata</i>
			<i>Rhaucus quinquelineatus</i>
			<i>Rhaucus vulneratus</i>
		Fissiphalliidae	<i>Fissiphallius sturmi</i>
			<i>Fissiphallius sympatricus</i>
		Stygnidae	<i>Phareus raptator</i>
	Cuenca alta del Río Cali	Cranaidae	<i>Holocranaus calus</i>
	Rio Meléndez	Cranaidae	<i>Holocranaus longipes</i>
	Río Nare	Cranaidae	<i>Holocranaus calcar</i>
Regional Protective Forest Reserve (RPFR)	Rios Blanco y Negro	Cosmetidae	<i>Libitia bipunctata</i>
		Stygnidae	<i>Phareus raptator</i>
Regional Protective Forest Reserve (RPFR)	Laguna de pantano redondo y el nacimiento Río Susagua	Cosmetidae	<i>Libitia bipunctata</i>
	Montes de Oca	Cosmetidae	<i>Eucynorta quadripustulata</i>
Flora and Fauna Sanctuary (FFS)	Guanentá-alto río Fonce	Cosmetidae	<i>Eulibitia clytemnestra</i>
	Iguaque	Cosmetidae	<i>Eulibitia maculata</i>
			<i>Libitia iguaque</i>
		Stygnidae	<i>Jabbastygnus huttonum</i>



In the species list, we utilize two types of dashes: the n-dash (–), used to indicate the reference of the original description, and the m-dash (—) to introduce each new subsequent bibliographic reference following the original description. After each species name, the m-dash is placed to separate a new citation.

Natural protected areas abbreviations: Civil Society Nature Reserve (**CSNR**); Flora and Fauna Sanctuary (**FFS**); National Protective Forest Reserve (**NPFR**); Regional Natural Park (**RNP**); Regional Protective Forest Reserve (**RPFR**); National Natural Parks (**NNP**); Regional Integrated Management Districts (**RDIM**).

## Results

### Species descriptions

**Order Opiliones Sundevall, 1833**

**Suborder Laniatores Thorell, 1876**

**Family Cranidae Roewer, 1913**

#### *Phalangodus* Gervais, 1842

**Included species.** *Phalangodus anacosmetus* Gervais, 1842 (type species); *Phalangodus briareos* Villarreal & García, 2016; *Phalangodus andresi* sp. nov.; *Phalangodus cottus* Villarreal & García, 2016; *Phalangodus gyes* Villarreal & García, 2016; *Phalangodus kuryi* Villarreal & García, 2016; *Phalangodus palpiconus* (Roewer, 1943).

**Diagnosis.** See Villarreal and García (2016).

#### *Phalangodus andresi* sp. nov.


<https://zoobank.org/B451F03A-F855-4F5B-8E90-8BDAF7890A11>

Figs 1–4

**Material examined.** • **Holotype:** ♂ (ICN-Ao-1908), COLOMBIA, Cundinamarca, San Antonio del Tequendama, R.N. Los Tunos (4.562234, -74.314527); 2,250 m; 3 Jun. 2018; (A. García, S. Galvis leg.). • **Paratypes:** • 3 ♀♀ (ICN-Ao-1909, with one used for description), same data as the holotype; • 2 ♂♂, 1 ♀ (ICN-Ao-1003), COLOMBIA, Cundinamarca, San Antonio del Tequendama, R.N. Los Tunos; 28 Aug. 2006; (F. Borrero leg.).

**Diagnosis.** *Phalangodus andresi* sp. nov. can be distinguished from all other species of the genus except *P. palpiconus* by the (1) presence of conspicuous granulation of mesotergal areas I–IV, lateral borders of dorsal scutum, ocularium and posterior region of the carapace (Figs 1A, B, 3A, B); (2) small size (males DSL ~ 10.7–11.9 mm), except *P. kuryi* (9.0–11.4 mm). It can be distinguished from the latter species by the ornamentation of the pedipalpal femur (with a very large ventroproximal tubercle absent in *P. kuryi*) (Fig. 3D), and the presence of ornamentation in leg IV of the males (Fig. 3G–J) (absent in *P. kuryi*), as well as the interocular distance, height of the ocularium and presence of paired tubercles near each eye (absent in *P. kuryi*). From *P. palpiconus*, the most morphologically similar species, it is distinguished by the ornamentation of the femur IV of the males, having a subdistal large and curved spine and a short distal bifid tubercle (Figs 2E, F, 3H, I) on the prolateral face (absent in *P. palpiconus*); lack of a retrolateral subdistal





**Figure 1.** *Phalangodus andresi* sp. nov. Male holotype (ICN-Ao-1908) **A** habitus, dorsal view **B** lateral view. Female paratype (ICN-Ao-1909): **C** dorsal view **D** lateral view. Scale bars: 3 mm.

spine in the same segment (present in *P. palpiconus* (Hara et al. 2014: figs 4, 5)); the shape of the VP of the penis, more elongated and with more marked medial constrictions, and the more basal position of MS-A/B groups.

**Description. Measurements of body and appendages.** Holotype ♂ (ICN-Ao-1908): DSL = 10.8; CL = 4.5; AL = 6.3; CW = 5.6; AW = 7.4; IOD = 1.6; BaCh = 2.7; FePp = 2.1, PaPp = 1.1, TiPp = 1.6, TaPp = 1.3, ClPp = 1.1; FeL I: 6.0; FeL II = 11.4; FeL III = 8.9; FeL IV = 14.1; TiL I = 4.6; TiL II = 11.3; TiL III = 5.3; TiL IV = 8.2. • Paratypes: • (ICN-AO-1909, ICN-AO-1003, 4 ♀♀, min–max): DSL = 8.3–9.3; CL = 2.9–4.5; AL = 4.0–7.8; CW = 4.9–5.8; AW = 7.2–7.9; IOD = 1.4–2.0; BaCh = 1.4–1.9; FePp = 1.9, PaPp = 1.1, TiPp = 1.4, TaPp = 1.1, ClPp = 1.0; FeL I = 2.9–3.6; FeL II = 8.1–10.6; FeL III = 7.3–9.4; FeL IV = 8.9–14.4; TiL I = 3.8–4.5; TiL II =




7.9–9.5; TiL III = 4.9–5.7; TiL IV = 5.9–9.0. • (ICN-AO-1003; 2 ♂♂, min–max): DSL = 10.7–11.8; CL = 4.9–5.3; AL = 6.6–7.4; CW = 5.3–6.3; AW = 7.4–8.8; IOD = 1.9–2.1; BaCh = 2.6–3.5; FePp = 1.9–2.1, PaPp = 1.0–1.2, TiPp = 1.5–1.7, TaPp = 1.2–1.4, ClPp = 1.1; FeL I = 5.1–6.4; FeL II = 11.1–11.1; FeL III = 9.4–10.6; FeL IV = 12.2–16.8; TiL I = 4.3–5.0; TiL II = 10.2–10.7; TiL III = 5.6–9.3; TiL IV = 8.4–10.2.

**Male holotype (ICN-Ao-1908). Dorsum** (Figs 1A, B, 3A, B). DS outline type alpha, with bulge longitudinally asymmetric widest at scutal groove II, lateral borders with granules only on the middle region. Carapace with few granules on the anterolateral and posterior region. Ocularium high, without median depression, with a paramedian pair of sharp tubercles and granules close to the eyes. Integumentary dome of ozopore raised and conspicuous. Abdominal scutum well delimited, divided into four well-marked scutal areas; I divided into left and right halves by invasion of the scutal area II; I and II granulated, with a pair of conspicuous medial tubercles, one tubercle on each side; III with a pair of paramedian acuminate spines and densely granulated; IV divided, with a row of six or seven granules on each side. Posterior border of the DS slightly convex and with a row of granules. Free tergites I–III with a row of granules.


**Venter.** Coxa I with a row of large tubercles of different sizes; II longer than I and III, with two median rows of low tubercles, the anterior one more conspicuous; III densely covered with irregular rows of small tubercles and with the posterior border sigmoid; IV strongly backward, with a median row of tubercles in the medium area and lateral border, and small tubercles densely distributed. Stigmatic area with a row of small tubercles on posterior border and minute granules sparsely distributed. Stigmata large, oval, and oblique. Free sternites with a row of small granules.

**Chelicerae** (Figs 2A, 3C). Segment I with well-defined bulla, with dorsomesal tubercles and a row of three large tubercles in the ectal region. Segment II swol-



**Figure 2.** *Phalangodus andresi* sp. nov. Male holotype (ICN-Ao-1908) **A** chelicera, frontal view **B** right palp, femur, ectal view **D** right leg IV: femur distal portion and patella, prolateral view **E** ditto, retrolateral view **F** ditto, ventral view **C** female, paratype (ICN-Ao-1909), right palp, femur, ectal view. Scale bars: 1 mm.






**Figure 3.** *Phalangodus andresi* sp. nov. Male holotype (ICN-Ao-1908) **A** habitus, dorsal view **B** habitus, lateral view **C** chelicera, frontal view **D** right pedipalp: trochanter, femur, and patella, ectal view **E** ditto, tibia, tarsus and claw, dorsoectal view **F** ditto, mesal view **G** right leg IV, trochanter and femur, dorsal view **H** ditto, femur, distal portion, prolateral view **I** ditto, ventral view **J** right leg IV, patella and tibia, ventral view. Scale bars: 1 mm.

len. Fixed finger with a proximal narrow and low tooth, a large gap and three subdistal median teeth. Movable finger with a proximal wide and low tooth, one large tooth and the distal inner surface irregularly dentate. Mesal side of the base of the fixed finger and near the base of the movable finger with setiferous tubercles of different sizes.

**Pedipalps** (Figs 2B, C, 3D–F). Trochanter with a dorsal pair of paramedian tubercles. Ventrally with large bifid tubercle in distal portion. Femur slightly compressed, dorsally curved, and ventrally straight in lateral view, with dorsoectal distal row of granules, one dorsal, and one ventral row of large forward projected tubercles (the apicalmost of the ventral row bifid and thicker than the remaining), the ventrodistal portion unarmed. Mesal and ectal faces without large tubercles. Patella short, cylindrical, and curved, with a dorsal row of short tubercles and small dorsodistal granules. Tibia dorsally granulated; tibia mesal lili ( $3 > 1 > 2 = 4$ ), ectal liili ( $4 > 1 > 5 > 2 > 3$ ). Tarsus dorsally granulated, tibia with spines only in the distal portion, mesal li, ectal liil ( $3 > 1 > 5 > 4 > 2$ ). Claw proximally swollen.





**Figure 4.** *Phalangodus andresi* sp. nov. Male paratype (ICN-Ao-1003). Penis, apical portion **A** dorsal view **B** lateral view **C** ventral view. Scale bars: 0.25 mm.

**Legs** (Figs 1A, 2D–F, 3B, 3G–J). Coxa I and III smooth; II with one dorsal tubercle; IV with one dorsodistal domed large tubercle, and a row of large tubercles in the prolateral border. Trochanters I–III unarmed; II with a prolateral row of small tubercles; III with one retrodistal tubercle and one group of small prolateral tubercles; IV with small prolateral granules sparsely distributed and one row of small tubercles in the retrolateral border. Femora I–III straight and with longitudinal rows of granules; IV sub-straight, densely granulated, ventrally with one large subdistal tubercle hook-like shaped and one distal rounded trifold tubercle. Ratio Fe IV/DSL = 1.85. Patellae I–IV granulated. Tibiae I–V straight and densely granulated, unarmed. Metatarsi I to IV with rows of small granules, I–III unarmed, IV with a ventro-distal border pair of spines. Tarsi III and IV with subparallel smooth claws and tarsal process. Tarsal counts: 8(3)-8(3), 13(3)-14(3), 8-8, 8-7.

**Penis** (Fig. 4). VP subsquare, with proximal and distal portion delimited by a medial constriction, and lateral rounded lobes in the proximal portion; distal border slightly concave. Glans + Stylus columnar, glans with some folds at the base, stylus normally thickened, shorter than the glans, substraight; styler caps ring-shaped. MS-A/B groups indistinguishable from each other, located on the proximal portion of the VP, with an increase in setae and composed of three to four pairs; MS-C/D groups located in the distal portion of the VP, separated from MS-A/B by a small gap, the setae rearranged into two irregular rows, one



laterodistal and one mesodorsal. MS-E presumably present, since they are very small (as in other members of the genus), and were not observable under the magnification used.

**Coloration (in alcohol)** (Figs 1, 2). Carapace, DS borders, posterior region and ocularium reticulated in Moderate brown 58, on background Light yellowish brown 76 (in females, it is Dark brown 59 on background Light brown 57). Abdominal scutum Light yellowish brown 76 (in females, it is moderate orange 53), with the scutal areas Moderate brown 58 (in females, it is Strong brown 55). Free tergites Dark grayish yellowish brown 81. Pedipalps, chelicerae and trochanters Strong brown 55 reticulated; remaining segment of the legs Strong brown 55 reticulated with fine mottled Deep orange 51. Stigmatic area Strong brown 55. Tip of cheliceral teeth Deep reddish brown 41.

**Female (ICN-Ao-1909)** (Fig. 1C, D). Differing from male by: ocularium slightly narrower; carapace shorter; coda wider (DS outline type alpha-keyhole); tubercles of area III slightly higher. Chelicerae non-hypertelic, with movable finger thinner. Pedipalpal femur lower and thinner in lateral view. Stigmatic area shorter. Genital operculum wider. Trochanters III and IV narrower; femur IV thinner, without large ventral-subdistal tubercle with hook-like shape and without a ventrodistal rounded trifold tubercle. For color differences, see the color description of the male.

**Ovipositor.** Dorsal lobes (dl) and ventral lobes (vl) rounded, with four and two pairs of large, acuminate, single-tipped setae, respectively. The dl with three pairs of dorsal setae (ds) distally located and one pair basally located. Lateral region of the ovipositor with one pair of dorso-lateral setae (dls).

**Distribution.** Known only from the type locality.

**Etymology.** The species is named in honor of our colleague and friend, the arachnologist Andrés F. García, who has greatly enriched the field's knowledge of Opiliones in Colombia and described the vast majority of species within the genus *Phalangodus*; moreover, he was the collector of the type series for this species.

### **Ventrifurca Roewer, 1913**

**Included species.** *Ventrifurca abnormis* (Roewer, 1932); *Ventrifurca albipustulata* Roewer, 1913 (type species); *Ventrifurca caffeinica* Villarreal, Kury & Pinto-da-Rocha, 2015; *Ventrifurca dybasi* (Goodnight & Goodnight, 1947); *Ventrifurca phallaina* sp. nov.

**Diagnosis.** Villarreal et al. (2015).


#### ***Ventrifurca phallaina* sp. nov.**

<https://zoobank.org/8101A255-1326-46E3-91C7-5D564C59FFC1>

Figs 5–9

**Material examined.** • **Holotype:** ♂ (CIUQ-020631), COLOMBIA – Valle del Cauca, Buenaventura, (3.8375, -77.2436); 56 m; 12 Nov. 2022; (A. L. García, L. Delgado-Santa leg.). • **Paratypes:** • 1 ♀ (CIUQ-020632), same data as the holotype • 1 ♂ (MIZA 0105869) same data as the holotype • 1 ♂ (MNRJ 1596) same data as the holotype. **Other material examined.** • 3 ♂♂, 1 ♀ (CIUQ-020634) same data as the holotype.






**Figure 5.** *Ventrifurca phallaina* sp. nov. Male holotype (CIUQ-020631) **A** dorsal view **B** lateral view **C** lateral view **D** right chelicera, frontal view. Scale bars: 1 mm.

**Diagnosis.** *Ventrifurca phallaina* sp. nov. can be distinguished from *V. abnormis* by the scarce ornamentation of the mesotergal areas I to III; also, can be distinguished from all other species of the genus by the ornamentation pattern of the yellow tubercles on dorsal scutum, restricted to the lateral of carapace and posterolateral portion of DS (Figs 5A, B, 6A, B, 9A, B); the presence of a yellow spot on the anterolateral portion of the carapace; the shape and size of posteroventral projections on the stigmatic area, simple and short (Figs 5C, 6C), instead of bifid and globular (*V. albipustulata*) or large and digitiform or curve (*V. caffeinica* and *V. abnormis*); and the absence of yellow tubercles on the mesotergal area III, behind the paired spines (present in *V. albipustulata* and *V. caffeinica*). The genital morphology is very similar to *V. albipustulata*, with slight differences in relation to the shape of the concavity of the truncus (Fig. 8).





**Figure 6.** *Ventrifurca phallaina* sp. nov., male paratype (MIZA 0105869) **A** habitus, dorsal view **B** habitus, lateral view **C** stigmatic area, posterior portion, ventral view **D** chelicera, frontal view **E** right pedipalp, ectal view **F** ditto, mesal view **G** right leg: femur, dorsal view **H** ditto, patella and tibia, dorsal view. Scale bars: 1 mm.

**Description. Measurements of body and appendages.** Holotype: (CIUQ-020631 ♂). DSL = 8.3, CL = 3.1, AL:5.2, CW = 3.2, AW = 5.3, IOD = 0.9; BaCh = 0.8, FePp = 2.1, PaPp = 1.1, TiPp = 1.5, TaPp = 1.3, ClPp = 0.9; FeL I = 3.8, FeL II = 9.8, FeL III = 7.6, FeL IV = 9.8, TiL I = 2.8, TiL II = 7.4, TiL III = 3.0, TiL IV = 5.4. Paratypes: • (MNRJ 1596, 1 ♂; MIZA 0105869 1 ♂): DSL = 7.8–8.4; CL = 3.4–3.2, AL = 3.9–5.3, CW = 4.0–4.3, AW = 6.5–6.7, IOD = 1.1–1.2; BaCh = 0.8–0.9; FePp = 2.3, PaPp = 1.3, TiPp = 1.7, TaPp = 1.2; ClPp = 1.3; FeL I = 4.6–4.8, FeL II = 10.6–10.8, FeL III = 7.4–8.1, FeL IV = 10.7–11.2, TiL I = 3.1–3.3, TiL II = 7.9–8.9, TiL III = 3.7–4.3, TiL IV = 5.9–6.0. • (CIUQ-020632, ♀): DSL = 6.0, CL = 2.7, AL = 2.9, CW = 3.0, AW = 5.2, IOD = 1.1; BaCh = 0.9; FePp = 2.2, PaPp = 1.2, TiPp = 1.6, TaPp = 1.1, ClPp = 1.1; FeL I = 2.6, FeL II = 7.4, FeL III = 6.2; FeL IV = 8.1, TiL I = 1.5, TiL II = 5.2, TiL III = 3.5, TiL IV = 4.6.

**Male holotype (CIUQ-020631). Dorsum** (Figs 5A, B, 6A, B). DS outline type Gamma pyriform. Mesotergum widest at groove III level; lateral margin of DS



with yellow rounded tubercles from carapace to area I, following outline of scutum. Carapace with three or four tubercles on each side of the anterior border. Ocularium low, armed with a paramedian pair of acuminate forward inclined tubercles and some small tubercles near the eyes. Integumentary dome of ozopore raised and conspicuous. Mesotergum well delimited, divided into three well-marked areas: area I divided into two roughly ellipsoidal halves by longitudinal groove, with a paramedian pair of large tubercles and four small tubercles; II penetrating subtly into I, with a median transverse row of eight small tubercles; III with a pair of paramedian acuminate high subparallel spines, located near the posterior margin; vestigial groove between III and IV present, partially blurred and located just behind the spines. Posterior border of DS straight. Free tergites I–III with a pair of paramedian large granules, I to II with one to three smaller laterodistal granules on each side; III armed with a posterior row of 11 tubercles, the paramedian pair much stouter.


**Venter** (Figs 5B, C, 6B, C). Coxa I with a row of tubercles of different size; coxa II longer than coxae I and III, with a cluster of distal tubercles of different size; coxa IV strongly backward. Stigmatic area with minute granules sparsely distributed, posterior border with posterodistal process very small. Stigmata large and oval.

**Chelicerae** (Figs 5D, 6D). Segment I with well-defined bulla, with a row of four or five tubercles in ectal region. Segment II slightly swollen, with a row of large



**Figure 7.** *Ventrifurca phallaina* sp. nov., male paratype (MIZA 0105869), right pedipalp **A** dorsoectal view **B** patella, tibia, tarsus, and claw, dorsomesal view **C** ectal view. Scale bars: 1 mm.





**Figure 8.** *Ventrifurca phallaina* sp. nov., male paratype (CIUQ-020633), penis **A** dorsal view **B** lateral view **C** ventral view. Abbreviations: MS = macrosetae of penis; MS-A = proximal group of setae; MS-B ventroproximal group of setae; MS-C = distal group of setae; MS-D = dorso medial group of setae; MS-E = ventrodistal group of reduced setae, SC = stylar caps. Figures not scaled.

setiferous tubercles near to the base of the movable finger and a few mesal setiferous tubercles. Fixed finger with three medial teeth, decreasing in size. Movable finger with one large and wide medial tooth, accompanied by two flat, low teeth on each side.

**Pedipalps** (Figs 6E, F, 7). Trochanter with tubercles, in dorsal and ventral region. Femur cylindrical, slightly elongated, armed with a longitudinal row of four tubercles in dorsal side and three tubercles in ventral side. Patella short, cylindrical, and curved, with row of tubercles in dorsal view, one ectodistal and two parallel rows of tubercles in mesal view tubercles.

**Legs** (Figs 5A–C, 6A, B, G, H). Coxae I and II with one anterior row of large tubercles; III with two rows and IV with two diagonal rows. Trochanter I with two prolateral, two retrolateral and four ventral tubercles; II with two prolateral, two retrolateral and five ventral tubercles; III with four prolateral, two retrolateral and four ventral tubercles; IV with one prolateral, one retrolateral and five ventral tubercles, more conspicuous than those from the other trochanters. Tubercles of trochanter IV more conspicuous than the others. Femora I to III straight, with longitudinal rows of granules; IV curved in its proximal portion, with seven longitudinal rows of tubercles, the retrolateral row with large proximal tubercles, and one large subdistal tubercle on the same row. Patellae I–IV with small tubercles. Tibiae and metatarsi I–V straight and granulate, unarmed. Claws III and IV smooth, with two minute proximal ventral protuberances. Ratio Fe IV/DSL = 1.35. Tarsal process present. Tarsal counts 7(3)-7(3), 14(3)-14(3), 7-7, 7-7.

**Penis** (Fig. 8). Ventral plate (VP) with concave lateral margins, with subdistal constriction and the distal corners ear-shaped, apical wide u-shaped cleft. Glans + Stylus complex surpassing the VP, glans columnar, with folds at the





Figure 9. *Ventrifurca phallaina* sp. nov., female paratype (CIUQ-020632) **A** dorsal view **B** lateral view **C** ventral view. Scale bars: 1 mm.

base, stylus normally thickened, sub-straight; styler caps foliar-shaped, short. MS-A1–A2 straight, located at the distal portion of the VP; MS-B absent; MS-C1–C3 curve, close to each other, and located in the distal part of the VP; MS-D1 larger than MS-C group, curve and located medially to MS-C, MS-D2 absent; MS-E1–E2 small, located on the flange of the VP. Truncus with ventral concavity next to the VP.

**Coloration (in alcohol)** (Fig. 5). Carapace, border of the DS, scutal grooves and adjacencies, and free tergites reticulated Dark brown 59, on background Brownish orange 54 and Brilliant orange yellow 67. Scutal areas Brilliant yellow 83. Trochanters I–III same as the carapace; IV Blackish red 21 and Brownish black 65. Pedipalps and chelicerae strong yellow 84 reticulated. Stigmatic area Strong yellowish brown 74. Tip of cheliceral teeth Dark olive 108.

**Female (CIUQ-020632)** (Fig. 9). Differing from male by carapace shorter than male, coda wider, femur IV thinner and with less conspicuous spination, free tergite spines larger, and stigmatic area without posterior process.

**Distribution.** Known only from the type locality.

**Etymology.** *Phallaina* is a Greek word that means whale. It is used in apposition as a reference to the humpbacked shape of the species in lateral view.



## Family Ampycidae Kury, 2003

### *Ampycella* Roewer, 1929

**Included species.** *Ampycella frizzellae* (Mello-Leitão, 1942); *Ampycella spiniventris* Roewer, 1929 (type species); *Ampycella fortunata* sp. nov.

**Diagnosis.** Outline of DS type alpha; four mesotergal areas well defined and unarmed; DS with small yellowish tubercle on the lateral margins (Fig. 10A, B, F). Free tergites II and III with a posteromedial spine (Fig. 11B). Penis (only known to *A. fortunata* sp. nov.): VP sub-rectangular with medial constriction, giving it a guitar-shaped appearance, and distal border without cleft; MS-A1–A3 in the medial portion of the VP; MS-B absent; MS-C1–C3 distally located, similar to each other; MS-D1 as large as MS-A similar to each other, medially located and pointing distally. Stylus clearly differentiated from the glans, slightly curved in the distal portion (Fig. 12).

### *Ampycella fortunata* sp. nov.

<https://zoobank.org/89231F4C-833A-4627-A866-2F33D9849AE0>

Figs 10–12


**Material examined.** • **Holotype:** ♂ (CIUQ-020635), COLOMBIA – Valle del Cauca, Buenaventura, (3.8375, -77.2436); 56 m; 12 Nov. 2022; (A. L. García, L. Delgado-Santa leg.). • **Paratypes:** • 1 ♀ (CIUQ-020636), same data as the holotype • 2 ♂♂ (CIUQ-020637; CIUQ-020638) same data as the holotype.

**Diagnosis.** *Ampycella fortunata* sp. nov. can be distinguished from the other two species of the genus by the size ratio between the spines on the free tergites: spine on free tergite II larger than the spine on free tergite III (Figs 10A, B, F, 11A, B) (vs III larger than II). Additionally, it can be distinguished from *A. frizzellae* by having a low ocularium with two small tubercles (vs high ocularium with two erect spines) and the entire scutal area II (vs divided into three lobes); and from *A. spiniventris* for having the anterolateral part of scutal area II projected forwards, anterior to the middle part (vs anterolateral parts similar to the medial portion), and having longer dorsal spines on the coxa IV.

**Description. Measurements of body and appendages.** Holotype: (CIUQ-020635) DSL = 3.5, CL = 1.5, AL = 1.6, CW = 1.6, AW = 3.2, IOD = 0.4, BaCh = 0.5, FePp = 1.1; PaPp = 0.6, TiPp = 0.6, TaPp = 0.5, ClPp = 0.3, FeL I = 1.7, FeL II = 2.8, FeL III = 2.2, FeL IV = 2.4, TiL I = 1.3, TiL II = 2.2, TiL III = 1.4, TiL IV = 1.4. Paratype: (CIUQ-020636 ♀) DSL = 3.5, CL = 1.4, AL = 2.0, CW = 1.7, AW = 3.4, IOD = 0.4, BaCh = 0.5, FePp = 1.1; PaPp = 0.6, TiPp = 0.7, TaPp = 0.5; ClPp = 0.3, Fe I = 1.4, FeL II = 2.7, FeL III = 2.1, FeL IV = 2.4, TiL I = 1.2, TiL II = 2.2, TiL III = 1.3, TiL IV = 1.5. Paratypes: (CIUQ-020637; CIUQ-020638, 2 ♂♂, min–max) DSL = 3.3–3.4, CL = 1.4–1.4, AL = 1.6–1.6, CW = 1.6–1.6, AW = 3.2–3.2, IOD = 0.4–0.4, BaCh = 0.4–0.4, FePp = 1.1–1.1; PaPp = 0.5–0.6, TiPp = 0.5–0.6, TaPp = 0.5–0.6, ClPp = 0.3–0.3, FeL I = 1.3–1.4, FeL II = 2.5–2.6, FeL III = 2.1–2.1, FeL IV = 2.2–2.3, TiL I = 1.1–1.1, TiL II = 2.1–2.1, TiL III = 1.2–1.3, TiL IV = 1.3–1.4.

**Male holotype (CIUQ-020635). Dorsum** (Figs 10A, B, 11A, B). DS outline type alpha. Dorsal scutum widest at scutal area II; lateral borders of dorsal scutum with yellowish granules only on the middle region. Carapace mostly smooth,






**Figure 10.** *Ampycella fortunata* sp. nov., male holotype (CIUQ-020635) **A** dorsal view **B** lateral view **C** ventral view **D** right pedipalp: mesal view **E** ditto, ectal view. Female paratype (CIUQ-020636): **F** dorsal view. Scale bars: 1 mm.

with a group of granules on the posterior region. Ocularium located slightly posterior to the ozopores, with a paramedian pair of small tubercles. Mesotergum well delimited, divided into four well-marked areas: I divided medially into left and right halves, with a medial large tubercle on each half and one small lateral tubercle and two anterior granules on each side; II and III entire, with a paramedian pair of large tubercles and two to three pairs on each side; IV divided into two halves, with four tubercles on each side. Posterior border of the DS substraight, with a row of tubercles. Free tergites I with a row of tubercles; II and III with a medial large spine (spine in II larger than in III) and some lateral tubercles, some of them yellowish colored.

**Venter** (Figs 10B, C, 11B). Stigmatic area with the posterior border convex, unarmed, only with minute granules sparsely distributed. Stigmata large, oval, and subparallel. Coxa I with two rows of large tubercles of different sizes; coxa II longer than coxae I and III, II and III with two rows of low tubercles, the anterior ones more conspicuous; coxa IV strongly backward and widened, conspicuously wider than the anterior coxae, with large tubercles in the posterior and lateral regions. Free sternites each with a row of small tubercles, the lateral larger.

**Chelicerae** (Fig. 11A). Chelicerae not swollen. Fixed finger with five teeth. Movable finger with ten teeth, one large tooth and distal inner surface dentate.





**Figure 11.** *Ampycella fortunata* sp. nov., male holotype (CIUQ-020635) **A** habitus, dorsal view **B** habitus, lateral view **C–F** tarsus I–IV, retrolateral view **G** right trochanter and femur IV, dorsal view **H** patella and tibia IV, dorsal view. Scale bars: 1 mm.


Mesal side of the base of the fixed finger and near the base of the movable finger with setiferous tubercles of different sizes.

**Pedipalps** (Figs 10D, E). Trochanter unarmed. Femur slightly compressed, sub straight in lateral view, unarmed. Patella with one large dorsodistal tubercle and some granules. Tibia shorter than femur, dorsally smooth; tibia mesal II, ectal III. Tarsus shorter than tibia, dorsally smooth; tarsus mesal III, ectal lili.

**Legs** (Figs 10A, 11C–H). Coxa I to III with one anterior row of small tubercles; IV with five longitudinal rows and with a very large dorsodistal spiniform tubercle. Trochanters I and II with two prolateral, two retrolateral, and three ventral tubercles; III with three prolateral, three retrolateral, and three ventral tubercles; IV with three prolateral, three retrolateral, and four ventral tubercles. Tubercles of trochanter IV are more conspicuous than the others. Femora I and II without conspicuous ornamentation; femora III and IV with seven rows of spines. Patellae I–IV with small tubercles. Tibiae I–IV straight and with tubercles; IV with a row of conspicuous tubercles. Claws III and IV smooth. Ratio Fe IV/DSL = 0.69. Tarsal counts 5(3)-5(3), 10(3)-10(3), 6-6, 6-6.

**Penis** (Fig. 12). Ventral plate (VP) sub-rectangular with medial constriction giving it a guitar-shaped appearance; distal border straight, without distal cleft. Glans + Stylus complex columnar, both parts well differentiated; stylus normally thickened, slightly curved S-shaped. MS-A1–A3 aligned in the basal part of





**Figure 12.** *Ampycella fortunata* sp. nov., male paratype (CIUQ-020637), penis **A, B** lateral view **C, D** ventral view **E, F** dorsolateral view. Figures not scaled.

the VP; MS-B absent; MS-C1–C3 aligned distally, similar in size and shape; MS-D1 large, similar to MS-A, distally pointed, MS-D2 absent; MS-E1-E2 very small, near MS-C cluster.

**Coloration (in alcohol)** (Fig. 10A–E). Carapace, chelicerae, and legs reticulated Dark yellowish brown (78) on Strong yellowish brown (74), borders of the DS and free tergites Dark yellowish brown (78). Abdominal scutum Strong yellowish brown (74), with the scutal areas Dark yellowish brown (78); Coxae dorsally reticulated on background Dark orange yellow (72). Pedipalps, reticulated. Stigmatic area with two oval areas Dark orange yellow (72) surrounded by Dark yellowish brown (78). Tarsomeri III and IV Grayish olive (110).

**Female (CIUQ-020636)** (Fig. 10F). Distinguished from male by carapace broader and the coda wider, genital operculum wider, spines on coxae IV present but less conspicuous than those found in males. and distal spine of femur smaller than in males.



**Distribution.** Known only from the type locality.

**Etymology.** *fortunata* is a Latin word that means happy, lucky, and blessed. It is used in reference to the meaning of the name of the type locality Buenaventura.

**Remarks.** Among the known genitalia, only *Licornus tama* Villarreal & Kury, 2012 and *Hernandarioides plana* F.O. Pickard-Cambridge, 1905 have an elongated ventral plate. However, both genera have a subtle distal cleft in the distal border of the VP, which is not the typical deep cleft seen in the subfamily (e.g., *Hexabunus* Roewer, 1913, *Hutamaia* Soares & Soares, 1977, *Thaumatochranaus* Roewer, 1932). *Ampycella fortunata* sp. nov. shares the presence of an elongated ventral plate, but it is unique in not having a distal cleft. Additionally, a pair of large, apically pointing MS-D is observed in *Ampycella fortunata*, which is also observed in some of the genera whose genitalia is known (e.g., *Hexabunus*, *Hernandarioides* Pickard-Cambridge, 1905).

## Faunistics

### Species list recorded or described from Colombia

Only those species described, registered, or modified after 2003 are provided with a detailed logonymy here. For the remaining species, refer to the catalog by Kury (2003). When possible, we added information about the WWF ecoregion where the species is found according to available records.

#### Suborder Laniatores Thorell, 1876

#### Infraorder Grassatores Kury, 2002

#### Incertae sedis

#### *Cleombrotus minutus* Sørensen, 1932

**Distribution.** Unknown department.

#### *Philacarus samoides* Sørensen, 1932

**Distribution.** Unknown department. Originally cited from Ecuador, but Kury (2003) commented: “Neu Granada” (this name refers to the region of Darien Gulf, bordering Panama).

**Remarks.** Nueva Granada was a historical region of northern South America. The Viceroyalty of New Granada (1717–1723, 1739–1810, 1815–1822) was a political and administrative entity established by Spain in the 18<sup>th</sup> century that encompassed a large part of the Andean region of South America, including present-day Colombia, Ecuador, Panama, and Venezuela. It was governed by a viceroy appointed by the Spanish crown. After gaining independence from Spain in 1819, the Republic of New Granada (1831–1858) was established as a sovereign state in the northern Andean region of South America, encompassing present-day mainly Colombia and Panama. The republic underwent various changes in its name and territorial boundaries.



**Gonyleptoidea Sundevall, 1833**  
**Agoristenidae Šilhavý, 1973**

***Andrescava sturmi* Roewer, 1963**

**Distribution.** Huila. Magdalena Valley dry forests (NT0221).

***Avima scabra* (Roewer, 1963)**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Avima troglobia* (Pinto-da-Rocha, 1996)**

*Trinella troglobia* Pinto-da-Rocha, 1996: 321, figs 4, 8, 15, 16. — Kury 2003: 34.  
*Avima troglobia* — Villarreal and Kury 2009: 3. — García et al. (2022): 59, figs 4, 6A.

**Distribution.** La Guajira. Cordillera Oriental montane forests (NT0118).

**Remarks.** The type locality for this species is the state of Zulia, in Venezuela. It was recorded from Colombia by García et al. (2022).

***Avima tuttifrutti* García & Pastrana-M, 2021**

*Avima tuttifrutti* García & Pastrana-M, 2021: 2, figs 1–4.

**Distribution.** Córdoba. Northern Andean montane forest (NT0145).

***Avima venezuelica* Soares & Avram, 1981**

*Avima venezuelica* Soares & Avram, 1981: 95. — Villarreal and Kury 2009: 67. — García et al. (2022): 60, figs 5, 6A.

*Vima venezuelica* — González-Sponga 1987: 543, figs 708–713.

*Trinella venezuelica* — Pinto-da-Rocha 1996: 323. — Kury 2003: 34.

**Distribution.** La Guajira. Cordillera Oriental montane forests (NT0118).

**Remarks.** The type locality for this species is the state of Zulia, in Venezuela. It was recorded from Colombia by García et al. (2022).

***Avima wayuunaiki* García, Vargas & Gutiérrez, 2022**

*Avima wayuunaiki* García et al., 2022: 56, figs 1–3, 6A, C.

**Distribution.** La Guajira. Guajira-Barranquilla xeric scrub (NT1308).



***Barinas guanenta* García & Ahumada-C., 2022**

*Barinas guanenta* García & Ahumada-C., 2022: 3, figs 1, 2, 4.

**Distribution.** Santander. Magdalena Valley montane forests (NT0136).

***Barinas piragua* Ahumada-C. & García, 2020**

*Barinas piragua* Ahumada-C. & García in Ahumada-C. et al., 2020: 635, figs 2–5.  
— García and Ahumada-C. 2022: 5.

**Distribution.** Bolívar, La Guajira, Magdalena. Magdalena-Urabá moist forests (NT0137), Guajira-Barranquilla xeric scrub (NT1308), and Amazon-Orinoco-Southern Caribbean mangroves (NT1401).

***Leptostygnus leptochirus* Mello-Leitão, 1940**

*Leptostygnus leptochirus* Mello-Leitão, 1940: 306. — Soares 1945: 383. — Soares et al. 1992: 7. — Kury 1993: 129, figs 1–6. — Flórez and Sánchez-C 1995: 369. — García and Villarreal 2020: 5, figs 1, 8, 9.

**Distribution.** Norte de Santander. Catatumbo moist forests (NT0108).

***Leptostygnus yarigui* García & Villarreal, 2020**

*Leptostygnus yarigui* García & Villarreal, 2020: 7, figs 3–6, 8, 9.

**Distribution.** Santander. Magdalena Valley montane forests (NT0136).

***Muscopilio onod* Villarreal & García, 2021**

*Muscopilio onod* Villarreal & García, 2021: 157, figs 1–10.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Nemastygnus ovalis* Roewer, 1929**

*Nemastygnus ovalis* Roewer, 1929: 277, fig. 44. — Kury 2003: 145. — Pinto-da-Rocha et al. 2012: 30, figs 1A–C.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).



***Paravima lokura* García & Villarreal, 2023**

*Paravima lokura* García & Villarreal, 2023: 420, figs 5, 6, 18.

**Distribution.** Norte de Santander. Cordillera Oriental montane forests (NT0118).

***Paravima magistri* García & Villarreal, 2023**

*Paravima magistri* García & Villarreal, 2023: 424, figs 7, 8, 18, 19G.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Sabanilla ornata* Roewer, 1913**

**Distribution.** Atlántico. Guajira-Barranquilla xeric scrub (NT1308).

***Vima panita* García & Kury, 2020**

*Vima panita* García & Kury, 2020: 71, figs 5–8.

**Distribution.** Caquetá. Napo moist forests (NT0142)

**Ampycidae Kury, 2003**

***Ampycella fortunata* sp. nov.**

**Distribution.** Valle del Cauca. Chocó-Darién moist forests (NT0115).

***Ampycus telifer* (Butler, 1873)**

*Gonyleptes telifer* Butler, 1873: 116, figs 3, 4.

*Itequahy ensiferus* Mello-Leitão, 1949: 29, fig. 9.

*Ampycus telifer* — Simon 1879: 241. — Roewer 1913: 49, fig. 18; 1923: 411, fig. 506. — Mello-Leitão 1923: 117; 1932: 208, fig. 119. — Goodnight and Goodnight 1943: 10. — Soares and Soares 1954: 234. — García 2014: 93, fig. 1a, b.

**Distribution.** Amazonas. Solimões-Japurá moist forests (NT0163).

**Remarks.** The type locality of this species is in the state of Amazonas, in Brazil. It was recorded for Colombia by García (2014).

***Licornus tama* Villarreal & Kury, 2012**

*Licornus tama* Villarreal & Kury, 2012: 73, figs 1–19. — García 2014: 93, fig. 2a, b.



**Distribution.** Meta. Apure-Villavicencio dry forests (NT0201).

**Remarks.** The type locality for this species is the state of Táchira, in Venezuela. Recorded for Colombia by García (2014).

***Thaumatochranaus magnificus* Hara, Bragagnolo & Pinto-da-Rocha, 2017**

*Thaumatochranaus magnificus* Hara et al., 2017: 463, figs 3, 4, 7D, E.

**Distribution.** Amazonas. Caqueta moist forests (NT0107).

***Thaumatochranaus splendidus* Hara, Bragagnolo & Pinto-da-Rocha, 2017**

*Thaumatochranaus splendidus* Hara et al., 2017: 466, figs 5, 6, 7F, G.

**Distribution.** Amazonas. Solimões-Japurá moist forests (NT0163).

**Cosmetidae Koch, 1839**

***Cosmetus acanti* Medrano, Kury & Martínez, 2021**

*Cosmetus acanti* Medrano et al., 2021b: 435, figs 5A–D, 6A–E, 7A–C, 11.

**Distribution.** Chocó. Chocó-Darién moist forests (NT0115).

***Cosmetus villarreali* Medrano, Kury & Martínez, 2021**

*Cosmetus villarreali* Medrano et al., 2021b: 439, figs 8A–D, 9A–H, 10A–C, 11.

**Distribution.** Chocó. Chocó-Darién moist forests (NT0115).

***Cynorta calcarapicalis* Roewer, 1912**

**Distribution.** Bolívar (type locality) and Cundinamarca. Sinú Valley dry forests (NT0229) and Magdalena Valley montane forests (NT0136).

***Cynorta clavipus* Roewer, 1928**

**Distribution.** Unknown department.

***Cynorta clunipecten* Roewer, 1947**

**Distribution.** Unknown department.



**Remarks.** Kury (2003) refers to the Boyacá or Santander departments as possible localities.

***Cynorta dariensis* Roewer, 1925**

**Distribution.** Chocó. Chocó-Darién moist forests (NT0115).

***Cynorta lateralis* Roewer, 1928**

**Distribution.** Chocó. Chocó-Darién moist forests (NT0115).

***Cynorta maculorum* Goodnight & Goodnight, 1943**

**Distribution.** Unknown department.

***Cynorta rorida* Roewer, 1928**

**Distribution.** Chocó. Chocó-Darién moist forests (NT0115).

***Cynorta simplex* Roewer, 1928**

**Distribution.** Unknown department.

***Cynortellina lineata* Roewer, 1915**

**Distribution.** Tolima (without a specific locality).

***Cynortosoma reticulatum* Roewer, 1947**

**Distribution.** Tolima (without a specific locality).

***Erginulus australis* (Roewer, 1916)**

**Distribution.** Unknown department.

***Eucynorta quadripustulata* (Simon, 1879)**

*Cynorta quadripustulata* Simon, 1879: 196.

*Eucynorta quadripustulata* — Roewer 1912: 55. — Ahumada-C. et al. 2022: 2, figs 1–4, 5D, 6.



**Distribution.** Atlántico, Bolívar, Cesar, La Guajira, Magdalena. Amazon-Orinoco-Southern Caribbean mangroves (NT1401), Guajira-Barranquilla xeric scrub (NT1308), Magdalena-Urabá moist forests (NT0137), Cordillera Oriental montane forests (NT0118) and Sinú Valley dry forests (NT0229).

***Eucynorta venosa* Roewer, 1928**

**Distribution.** Chocó. Chocó-Darién moist forests (NT0115).

***Eucynortella orbicularis* Roewer, 1947**

**Distribution.** Unknown department.

**Remarks.** Flórez and Sánchez-C (1995) refers to the Boyacá or Santander departments as possible localities.

***Eucynortula ypsilon* Roewer, 1925**

**Distribution.** Chocó. Chocó-Darién moist forests (NT0115).

***Eulibitia castor* Medrano & Kury, 2017**

*Eulibitia castor* Medrano & Kury, 2017: 8, figs 5–7, 34.

**Distribution.** Boyacá. Cordillera Oriental montane forests (NT0118).

***Eulibitia chacuamarei* Pinzón, Damron & Pinto-da-Rocha, 2021**

*Eulibitia chacuamarei* Pinzón et al., 2021: 205, figs 1, 2.

**Distribution.** Casanare. Llanos (NT0709).

***Eulibitia clytemnestra* Medrano & Kury, 2017**

*Eulibitia clytemnestra* Medrano & Kury, 2017: 12, figs 8–10, 33.

**Distribution.** Santander. Magdalena Valley montane forests (NT0136).

***Eulibitia ectroxantha* (Mello-Leitão, 1941)**

*Brachylibitia ectroxantha* Mello-Leitão, 1941: 166, fig. 1.

*Cynorta ectroxantha* — Goodnight and Goodnight 1953: 38.



*Platymessa ectroxantha* — Medrano and Kury 2016: 57. — Tagged as species inquirenda.

*Eulibitia ectroxantha* — Medrano and Kury 2017: 16, fig. 54. — Medrano et al. 2019: 346, figs 1–4.

**Distribution.** Boyacá. Northern Andean páramo (NT1006) and Magdalena Valley montane forests (NT0136).

### ***Eulibitia helena* Medrano & Kury, 2017**

*Eulibitia helena* Medrano & Kury, 2017: 17, figs 11–14, 34.

**Distribution.** Norte de Santander. Cordillera Oriental montane forests (NT0118).

### ***Eulibitia h-inscriptum* (Mello-Leitão, 1941)**

*Platymessa h-inscripta* Mello-Leitão, 1941: 167, fig. 2.

*Platymessa nigrolimbata* Mello-Leitão, 1941: 168, fig. 3. — B. Soares 1945: 344.

*Cynorta h-inscripta* — H. Soares 1970: 325.

*Platymessa h-inscriptum* — Kury and Alonso-Zarazaga 2011: 51. — Medrano and Kury 2016: 58, figs 1–29.

*Eulibitia h-inscriptum* — Medrano and Kury 2017: 17, fig. 32.

**Distribution.** Boyacá and Santander. Magdalena Valley montane forests (NT0136) and Northern Andean páramo (NT1006).

### ***Eulibitia leda* Medrano & Kury, 2017**

*Eulibitia leda* Medrano & Kury, 2017: 22, figs 15–18, 33.

**Distribution.** Santander. Magdalena Valley montane forests (NT0136).

### ***Eulibitia maculata* Roewer, 1912**

*Eulibitia maculata* Roewer, 1912: 17, pl. 1, fig. 1. — Roewer 1923: 298, fig. 320. — Sørensen in Henriksen 1932: 389. — B. Soares 1945: 343. — Weidner 1959: 122. — Pinto-da-Rocha and Hara 2011: 2. — Medrano and Kury 2017: 27, figs 3, 19–22, 32.

*Libitia (Messa) castanea* Sørensen in Henriksen, 1932: 415.

*Paramessa castanea* — Mello-Leitão 1933: 109.

*Messatana castanea* — Strand 1942: 398.

**Distribution.** Boyacá, Cundinamarca, and Tolima. Northern Andean páramo (NT1006) and Magdalena Valley montane forests (NT0136).



***Eulibitia pollux* Medrano & Kury, 2017**

*Eulibitia pollux* Medrano & Kury, 2017: 34, figs 23–26, 31B, 34.

**Distribution.** Boyacá. Magdalena Valley montane forests (NT0136).

***Eulibitia scalaris* (Sørensen, 1932)**

*Libitia* (*Messa*) *scalaris* Sørensen in Henriksen, 1932: 414.

*Acromares lateralis* Goodnight & Goodnight, 1943: 2, fig. 7. — Flórez and Sánchez-C 1995: 368.

*Messa scalaris* — Mello-Leitão 1933: 112.

*Messatana scalaris* — Strand 1942: 398.

*Eulibitia scalaris* — Kury and Medrano 2017: 40, figs 27–30, 31A, 33.

*Cynorta lateralis* — Goodnight and Goodnight 1953: 38.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Eulibitia victoriae* (Pinzón-M. & Townsend, 2017)**

*Platymessa victoriae* Pinzón-M. et al., 2017: 61, figs 1–7.

*Eulibitia victoriae* — Kury et al. 2020: 14.

**Distribution.** Cesar. Cordillera Oriental montane forests (NT0118).

***Flirtea alpha* (Sørensen, 1932)**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Flirtea fusca* (Sørensen, 1932)**

**Distribution.** Unknown department.

***Flirtea granulosa* (Simon, 1879)**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Flirtea limbata* (Sørensen, 1932)**

**Distribution.** Unknown department.



***Flirtea militaris* (Simon, 1879)**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Flirtea tuberculata* (Sørensen, 1932)**

**Distribution.** Unknown department.

***Flirtea ventricosa* (Simon, 1879)**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Libitia bipunctata* Sørensen, 1932**

*Libitia (Libitia) bipunctata* Sørensen in Henriksen, 1932: 417.

*Libitia bipunctata* — Mello-Leitão 1933: 109. — Medrano, et al. 2020: 5, figs 1–4, 14.

*Libitiella bipunctata* — Roewer 1947: 8, pl. 1, fig. 3.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136), Cordillera Oriental montane forests (NT0118) and Northern Andean páramo (NT1006).

***Libitia cordata* (Gervais, 1844)**

*Cosmetus cordatus* Gervais, 1844: 117, pl. 46, fig. 9.

*Libitia cordata* — Butler 1873: 115. — Simon 1879: 216. — Roewer 1912: 12; 1923: 293. — Mello-Leitão 1923: 108; 1932: 53; 1933: 109. — Medrano et al. 2020: 10, figs 5–8, 14.

*Libitia (Libitia) cordata* — Sørensen in Henriksen 1932: 419.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136) and Northern Andean páramo (NT1006).

***Libitia gandalf* Medrano, Ázara & Kury, 2020**

*Libitia gandalf* Medrano et al., 2020: 14, figs 9, 10, 14.

**Distribution.** Meta. Cordillera Oriental montane forests (NT0118).

***Libitia iguaque* Medrano, Ázara & Kury, 2020**

*Libitia iguaque* Medrano et al., 2020: 16, figs 11, 12, 14.

**Distribution.** Boyacá. Northern Andean páramo (NT1006).



***Metacynorta bella* Roewer, 1963**

**Distribution.** Putumayo. Napo moist forests (NT0142).

***Meterginus affinis* Roewer, 1963**

**Distribution.** Putumayo. Napo moist forests (NT0142).

***Meterginus flavicinctus* (Gervais, 1842)**

**Distribution.** Cundinamarca. Magdalena Valley dry forests (NT0221) and Magdalena Valley montane forests (NT0136).

***Meterginus marginellus* (Simon, 1879)**

**Distribution.** Unknown department.

***Meterginus marmoratus* (Roewer, 1912)**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Meterginus obscurus* (Sørensen, 1932)**

**Distribution.** Unknown department.

***Meterginus prosopis* Roewer, 1912**

**Distribution.** Cundinamarca, Meta, and Tolima. Magdalena Valley montane forests (NT0136) and Cordillera Oriental montane forests (NT0118).

***Meterginus simonis* (With, 1932)**

*Rhaucus (Erginus) simonis* With in Henriksen, 1932: 350.

*Meterginus simonis* — Mello-Leitão 1933: 110. — Kury 2003: 73. — Kury et al. 2020: 17.

**Distribution.** Unknown department.

***Meterginus togatus* (Sørensen, 1932)**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).



***Paecilaema altaspinulatum* Goodnight & Goodnight, 1943**

**Distribution.** Unknown department.

***Paecilaema atroluteum* Roewer, 1912**

**Distribution.** Atlántico. Guajira-Barranquilla xeric scrub (NT1308).

***Paecilaema contextum* Roewer, 1928**

**Distribution.** Unknown department.

***Paecilaema distinctum* Roewer, 1915**

**Distribution.** Atlántico. Guajira-Barranquilla xeric scrub (NT1308).

***Paraprotus atroluteus* Roewer, 1912**

**Distribution.** Atlántico. Guajira-Barranquilla xeric scrub (NT1308).

***Protus speciosus* (Roewer, 1927)**

*Paraprotus speciosus* Roewer, 1927: 628, pl. 1, fig. 11.

*Protus speciosus* — Medrano et al. 2021a: 52.

**Distribution.** Amazonas. Solimões-Japurá moist forests (NT0163).

**Remarks.** The type locality for this species is Maranhão state, in Brazil. It was recorded for Colombia by Medrano et al. (2021a).

***Rhaucoides nasa* Medrano, García & Kury, 2022**

*Rhaucoides nasa* Medrano et al., 2022: 196, figs 7–9, 22.

**Distribution.** Cauca. Magdalena Valley montane forests (NT0136).

***Rhaucoides riveti* Roewer, 1914**

*Rhaucoides riveti* Roewer, 1914: 125, pl. 13, fig. 3. — Roewer 1923: 306, fig. 333.

*Rhaucoides festae* Roewer, 1925: 3, pl. 5, fig. 1. — *Pararhauculus sulfureus* Mello-Leitão, 1939: 171. — Established as a junior subjective synonym of *Rhaucoides festae* Roewer, 1925 by Kury (2003).

*Cumbalia octomaculata* Roewer, 1963: 51, pl. 9, fig. 13.



**Distribution.** Nariño. Northwestern Andean montane forests (NT0145).

***Rhaucus florezi* García & Kury, 2017**

*Rhaucus florezi* García & Kury, 2017: 426, figs 1e, 15–17, 18e, 23.

**Distribution.** Boyacá. Magdalena Valley montane forests (NT0136).

***Rhaucus papilionaceus* (Simon, 1879)**

*Erginus papilionaceus* Simon, 1879: 205.

*Flirtea papilionacea* — Roewer 1912: 77. — Roewer 1923: 347. — Roewer 1927: 593.  
— Mello-Leitão 1932: 78.

*Rhaucus* (*Erginus*) *papilionaceus* — Henriksen 1932: 352.

*Rhaucus papilionaceus* — García and Ahumada-C. 2018: 201, figs 1–8.

**Distribution.** Santander. Magdalena Valley montane forests (NT0136).

***Rhaucus quinquelineatus* Simon, 1879**

*Rhaucus quinquelineatus* Simon, 1879: 215. — Sørensen in Henriksen 1932: 358. — García and Kury 2017: 413, figs 1d, 6–8, 20a, 22.

*Pararhaucus obscurus* Pickard-Cambridge, 1905: 572. — Roewer 1912: 102. — Roewer 1923: 378.

*Metarhaucus albilineatus* Roewer, 1912: 147, pl. 7, figs 5, 6. — Flórez and Sánchez-C 1995: 368.

*Rhaucus* (*Rhaucus*) *muticus* Sørensen, 1932: 360.

*Rhaucus* (*Rhaucus*) *tristis* — Sørensen 1932: 363, fig. 26.

*Flirtea quinquelineata* — Roewer 1912: 76. — Roewer 1923: 346, fig. 393. — Mello-Leitão 1923: 112. — Roewer 1927: 594. — Mello-Leitão 1932: 79, fig. 47. — B. Soares 1945: 343. — Roewer 1963: 58. — Flórez and Sánchez-C 1995: 368.

*Paecilaema obscurum* — Goodnight and Goodnight 1953: 54. — Kury 2003: 78.

*Flirtea mutica* — Mello-Leitão 1933: 110.

*Flirtea tristis* — Mello-Leitão 1933: 110.

**Distribution.** Boyacá and Cundinamarca. Northern Andean páramo (NT1006), Magdalena Valley montane forests (NT0136) and Cordillera Oriental montane forests (NT0118).

***Rhaucus robustus* (Mello-Leitão, 1941)**

*Megarhaucus robustus* Mello-Leitão, 1941: 169, fig. 4.

*Rhaucus robustus* — García & Kury, 2017: 422, figs 1c, 12–14, 18d, 23.



**Distribution.** Boyacá and Santander. Northern Andean páramo (NT1006) and Magdalena Valley montane forests (NT0136).

***Rhaucus serripes* (Simon, 1879)**

*Erginus serripes* Simon, 1879: 204.

*Metarhaucus fuscus* Pickard-Cambridge, 1905: 572.

*Metarhaucus reticulatus* Roewer, 1912: 145, pl. 7, fig. 4.

*Rhaucus* (*Rhaucus*) *geographicus* Sørensen, 1932: 369.

*Flirtea paucimaculata* Roewer, 1963: 58, fig. 24. — Flórez and Sánchez-C 1995.

*Flirtea serripes* — Roewer 1912: 77; 1923: 348, figs 396–397; 1927: 593. — Mello-Leitão 1932: 78.

*Rhaucus* (*Erginus*) *serripes* — Henriksen in Sørensen 1932: 352.

*Rhaucus serripes* — García and Kury 2017: 417, figs 1f, 9–11, 18c, 23.

*Erginus fuscus* — Roewer, 1912: 68.

*Metarhaucus fuscus* — Roewer 1923: 342; 1927: 588.

*Erginus reticulatus* — Roewer 1912: 68.

*Metarhaucus reticulatus* — Roewer 1923: 343, fig. 387; 1927: 588; 1959: 80. — Forcart 1961: 51. — Flórez and Sánchez-C 1995: 368.

*Flirtea geographica* — Mello-Leitão 1933: 110.

**Distribution.** Boyacá and Cundinamarca. Northwestern Andean montane forests (NT0145), Magdalena Valley montane forests (NT0136), Cordillera Oriental montane forests (NT0118) and Northern Andean páramo (NT1006).

***Rhaucus vulneratus* Simon, 1879**

*Rhaucus vulneratus* Simon, 1879: 213. Pickard-Cambridge 1905: 572. — Roewer 1912: 78. — Roewer 1923: 349, fig. 400. — Mello-Leitão 1923: 113. — Mello-Leitão 1932: 57, fig. 23. — Mello-Leitão 1933: 110. — García and Kury 2017: 405, figs 1a, b, 2–5, 18a, 19a, b, 22.

*Neorhaucus aurolineatus* Pickard-Cambridge, 1905: 572.

*Rhaucus* (*Rhaucus*) *vulneratus* — Sørensen 1932: 355.

*Raucus vulneratus* — González-Sponga 1992: 427.

*Neorhaucus aurolineatus* — Roewer 1912: 25. — Roewer 1923: 305.

**Distribution.** Boyacá and Cundinamarca. Magdalena Valley montane forests (NT0136) and Northern Andean páramo (NT1006).

***Sibambea cincta* (Perty, 1833)**

*Discosoma cincta* Perty, 1833: 209, pl. 40, fig. 6.

*Discosomaticus sturmi* Roewer, 1963: 58, fig. 25.

*Sibambea cincta* — Medrano et al. 2021a: 56.



**Distribution.** Amazonas and Vaupés. Purus várzea (NT0156), Iquitos várzea (NT0128), Solimões-Japurá moist forests (NT0163) and Caqueta moist forests (NT0107).

**Remarks.** The type locality for this species is the state of Bahia, in Brazil. It was recorded for Colombia by Medrano et al. (2021a).

***Taito insperatus* (Soares, 1970)**

*Cynorta insperata* Soares, 1970: 323, fig. 3. — Kury 2003: 45.

*Taito insperatus* — Kury and Barros 2014: 36, figs 1d, 6ae–ai, 17d–f, 28.

**Distribution.** Amazonas. Purus várzea (NT0156).

***Taito medinae* Kury & Barros, 2014**

*Taito medinae* Kury & Barros, 2014: 39, figs 1a, 17j–l, 26a–g, 35.

**Distribution.** Amazonas. Solimões-Japurá moist forests (NT0163).

***Taito oblongatus* (Roewer, 1928)**

*Cynortula oblongata* Roewer, 1928: 576, fig. 28. — Mello-Leitão 1932: 58. — Kury 2003: 52.

*Taito oblongatus* — Kury and Barros 2014: 40, figs 1e, 6a–l, 17a–c, 29.

**Distribution.** Meta. Cordillera Oriental montane forests (NT0118).

***Zaraxolia devians* (Sørensen, 1932)**

**Distribution.** Unknown department.

**Cranaidae Roewer, 1913**

***Allocranaus columbianus* Roewer, 1915**

**Distribution.** Quindío. Magdalena Valley montane forests (NT0136).

***Cranaus albipustulatus* Roewer, 1943**

**Distribution.** La Guajira. Guajira-Barranquilla xeric scrub (NT1308).



***Cranaus chlorogaster* (Gervais, 1844)**

**Distribution.** Unknown department.

***Cranaus cinnamomeus* (Gervais, 1844)**

**Distribution.** Unknown department.

***Deriacrus simoni* Roewer, 1932**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Holocranaus calcar* (Roewer, 1912)**

**Distribution.** Antioquia. Cauca Valley montane forests (NT0109) and Magdalena Valley montane forests (NT0136).

***Holocranaus calus* (Goodnight & Goodnight, 1944)**

**Distribution.** Valle del Cauca. Cauca Valley montane forests (NT0109).

***Holocranaus longipes* Roewer, 1913**

*Holocranaus longipes* Roewer, 1913: 400, fig. 158. — Roewer 1923: 556, fig. 695. — 1932: 293, fig. 9. — Soares and Soares 1948: 602.

*Phareicranaus giganteus* Roewer, 1932: 299. — Kury 2003: 93. — Pinto-da-Rocha and Bonaldo 2011: 27.

**Distribution.** Unknown department.

**Remarks.** The type locality for this species is the Aguacatal River, Colombia. The locality is not precise (Kury 2003).

***Holocranaus pectinitibialis* (Roewer, 1914)**

*Tolimaius pectinitibialis* Roewer, 1914: 125, figs 12–13. — Roewer 1923: 558, fig. 698.

*Holocranaus pectinitibialis* — Soares and Soares 1948: 603. — Kury 2003: 93.

**Distribution.** Tolima. Magdalena Valley montane forests (NT0136).



***Holocranaus simplex* Roewer, 1913**

**Distribution.** Unknown department.

***Homocranaus tetracalcar* Roewer, 1915**

**Distribution.** Quindío. Magdalena Valley montane forests (NT0136).

***Isocranaus gorgonae* Hirst, 1926**

**Distribution.** Nariño. Chocó-Darién moist forests (NT0115).

***Isocranaus obscurus* Roewer, 1915**

**Distribution.** Meta. Apure-Villavicencio dry forests (NT0201).

***Megacranaus areolatus* Roewer, 1932**

**Distribution.** Tolima.

***Megacranaus pygoplus* Roewer, 1913**

**Distribution.** Antioquia. Northern Andean páramo (NT1006).

***Metacranaus tricalcaris* Roewer, 1913**

**Distribution.** Antioquia. Cauca Valley montane forests (NT0109).

***Neocranaus albiconspersus* Roewer, 1913**

**Distribution.** Maracaibo (without further specific locality data).

**Remark.** Several localities are called Maracaibo in Colombia and Venezuela. Kury (2003) suggests that it could be a Venezuelan locality. The other species in the genus is from Ecuador, and a taxonomic revision is underway to better understand the distribution and composition of the genus. At the moment, it is impossible to associate the species with any of the localities.

***Neocranaus armatissimus* (Mello-Leitão, 1941)**

*Mitobatulina armatissima* Mello-Leitão, 1941: 170, fig. 5. — Soares and Soares 1949: 235.

*Neocranaus armatissimus* — Kury 2003: 95.

**Distribution.** Nariño, Napo moist forests (NT0142).

***Panalus robustus* Goodnight & Goodnight, 1947**

**Distribution.** Cauca. Cauca Valley montane forests (NT0109).

***Paracranaus crassipalpis* Roewer, 1913**

**Distribution.** Atlántico. Guajira-Barranquilla xeric scrub (NT1308).

***Peripa simplex* Roewer, 1932**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Phalangodus anacosmetus* Gervais, 1842**

*Phalangodus anacosmetus* Gervais, 1842: 3, pl. 4; 1844: 114, pl. 46, fig. 3. — Erichson 1845: 267. — Thorell 1877: 115. — Simon 1879: 241. — Roewer 1913: 138, fig. 63. — 1923: 446, fig. 562. — Kästner 1937: 300, fig. 364. — Soares and Soares 1954: 289. — Kury 1996: 178, figs 1–4. — Acosta 1996: 224. — Hara et al. 2014: 569. — Villarreal and García 2016: 5, figs 1–3, 16a, 18–19. *Allochranaus giganteus* Mello-Leitão, 1940: 307, fig. 8. — B. Soares 1945: 349. — Soares and Soares 1948: 587.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Phalangodus briareos* Villarreal & García, 2016**

*Phalangodus briareos* Villarreal & García, 2016: 9, figs 4–6, 7a–d, 18, 19. — Barriga et al. 2019: 117.

**Distribution.** Santander. Magdalena Valley montane forests (NT0136).

***Phalangodus andresi* sp. nov.**

**Distribution.** Cundinamarca. Chocó-Darién Moist Forests (NT0115).

***Phalangodus cottus* Villarreal & García, 2016**

*Phalangodus cottus* Villarreal & García, 2016: 15, figs 8a, b, 9, 10, 18.



**Distribution.** Meta. Apure-Villavicencio dry forests (NT0201).

***Phalangodus gyes* Villarreal & García, 2016**

*Phalangodus gyes* Villarreal & García, 2016: 21, figs 7e–f, 8c–d, 11, 12, 18.

**Distribution.** Tolima. Magdalena Valley montane forests (NT0136).

***Phalangodus kuryi* Villarreal & García, 2016**

*Phalangodus kuryi* Villarreal & García, 2016: 25, figs 13–15, 18.

**Distribution.** Magdalena. Guajira-Barranquilla xeric scrub (NT 1308).

***Phareicranaus albigranulatus* Roewer, 1913**

*Phareicranaus albigranulatus* Roewer, 1913: 404, fig. 160. — Kury 2003: 96. — Pinto-da-Rocha and Bonaldo 2011: 27.

**Distribution.** Tolima. Magdalena Valley montane forests (NT0136).

**Remarks.** Pinto-da-Rocha and Bonaldo (2011) question the generic placement of this species.

***Phareicranaus albigyratus* Roewer, 1932**

*Phareicranaus albigyratus* Roewer, 1932: 303. — Kury 2003: 96. — Pinto-da-Rocha and Bonaldo 2011: 6.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Phareicranaus angelicus* (Roewer, 1963)**

*Santinezia angelica* Roewer, 1963: 69. — Pinto-da-Rocha and Kury 2003: 182. — Kury 2003: 97.

*Phareicranaus angelicus* — Pinto-da-Rocha and Bonaldo 2011: 8, fig. 4B, C.

**Distribution.** Meta. Cordillera Oriental montane forests (NT0118).

***Phareicranaus spinulatus* (Goodnight & Goodnight, 1943)**

*Santinezia spinulata* Goodnight & Goodnight, 1943: 9, figs 26–28. — Pinto-da-Rocha and Kury 2003: 177, 197, 198, 202, 204, 205, fig. 51. — Kury 2003: 99.

*Phareicranaus spinulatus* — Pinto-da-Rocha and Bonaldo 2011: 26, fig. 18C, D.

**Distribution.** Unknown department.

***Sibundoxia scripta* Roewer, 1963**

**Distribution.** Putumayo. Eastern Cordillera real montane forests (NT0121).

***Stygnicranaus alessandroi* Orrico & Kury, 2009**

*Stygnicranaus alessandroi* Orrico & Kury, 2009: 474, figs 2–5.

**Distribution.** Valle del Cauca. Cauca Valley montane forests (NT0109).

***Stygnicranaus poncedeleoni* Orrico & Kury, 2009**

*Stygnicranaus poncedeleoni* Orrico & Kury, 2009: 476, figs 6–11.

**Distribution.** Valle del Cauca. Northwestern Andean montane forests (NT0146).

***Tetracranaus zilchi* Roewer, 1963**

**Distribution.** Valle del Cauca. Northwestern Andean montane forests (NT0146) and Magdalena Valley montane forests (NT0136).

***Ventrifurca albipustulata* Roewer, 1913**

*Ventrifurca albipustulata* Roewer, 1913: 383, figs 149–150. — Kury 2003: 100. — Villarreal et al. 2015: 6, figs 1, 5, 7, 8, 10, 11, 13.

*Microcranaus pustulatus* Roewer, 1913: 353, fig. 137. — Kury 2003: 93.

*Cayabeus perlatus* Roewer, 1932: 337, fig. 53. — Kury 2003: 91.

**Distribution.** Antioquia. Cauca Valley montane forests (NT0109).

***Ventrifurca caffeinica* Villarreal, Kury & Pinto-da-Rocha, 2015**

*Ventrifurca caffeinica* Villarreal et al., 2015: 11, figs 2, 4, 6, 12, 13.

**Distribution.** Quindío. Cauca Valley montane forests (NT0109).

***Ventrifurca dybasi* (Goodnight & Goodnight, 1947)**

*Rhopalocranaus dybasi* Goodnight & Goodnight, 1947: 40, fig. 20.

*Neocranaus dybasi* — Kury 2003: 95.



*Ventrifurca dybasi* — Villarreal et al. 2015: 14, fig. 3A, B, C.

**Distribution.** Valle del Cauca. Cauca Valley montane forests (NT0109).

**Remarks.** Generic assignment from *Neocranaus* by Villarreal et al. (2015).

*Ventrifurca phallaina* sp. nov.

**Distribution.** Valle del Cauca. Chocó-Darién moist forests (NT0115).

*Ventrisudis mira* Roewer, 1963

**Distribution.** Huila. Patía Valley dry forests (NT0225).

**Manaosbiidae** Roewer, 1943

*Camelianus fuhrmanni* Roewer, 1912

**Distribution.** Antioquia. Cauca Valley montane forests (NT0109) and Magdalena Valley montane forests (NT0136).

*Cucutacola nigra* Mello-Leitão, 1940

**Distribution.** Norte de Santander. Catatumbo moist forests (NT0108).

*Gonogotus areolatus* Roewer, 1943

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

*Rhopalocranaus apiculatus* Roewer, 1932

**Distribution.** Unknown department.

*Rhopalocranaus atroluteus* Roewer, 1913

**Distribution.** Tolima. Magdalena Valley montane forests (NT0136).

*Rhopalocranaus columbianus* (Roewer, 1963)

*Microcranaus columbianus* Roewer, 1963: 64, figs 34, 35.

*Rhoplocranaus columbianus* — Villarreal et al. 2015: 14, fig. 15.

**Distribution.** Cauca. Patía Valley dry forests (NT0225).

***Rhopalocranaus ypsilon* Roewer, 1913**

**Distribution.** Tolima. Magdalena Valley montane forests (NT0136).

***Semostrus tarsalis* Roewer, 1943**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

**Nomoclastidae Roewer, 1943**

***Calcosma barasana* Pinto-da-Rocha & Bragagnolo, 2017**

*Calcosma barasana* Pinto-da-Rocha & Bragagnolo, 2017: 103, figs 4B, 8B, 10B, 13A, B.

**Distribution.** Vaupés. Caqueta moist forests (NT0107).

***Calcosma gracillima* Roewer, 1932**

*Calcosma gracillima* Roewer, 1932: 331, fig. 47. — Soares and Soares 1948: 591. — Kury 2003: 91. — Pinto-da-Rocha and Bragagnolo 2017: 110, figs 1E, 4D, 8D, 10C, 13E, F.

**Distribution.** Amazonas. Solimões-Japurá moist forests (NT0163).

***Nomoclastes quasimodo* Pinto-da-Rocha, 1997**

*Nomoclastes quasimodo* Pinto-da-Rocha, 1997: 171, figs 1–6, 449, 450, 587, 592, 596. — Kury 2003: 228.

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Nomoclastes taedifer* Sørensen, 1932**

*Nomoclastes taedifer* Sørensen, 1932: 300, fig. 11. — Roewer 1943: 36. — Pinto-da-Rocha 1997: 171, figs 7, 8. — Kury 2003: 228.

**Distribution.** Department unknown.

***Paraphalangodus synacanthus* Roewer, 1915, new family assignment**

**Distribution.** Tolima (without a specific locality).

**Remark.** The type locality for this species refers to Páramo del Tolima, located in the department of Tolima, which comprises four páramo



complexes, namely Las Hermosas, Chilí Barragán, Nevado del Huila-Moras, and Los Nevados. Therefore, the specific location within the Tolima department cannot be determined without further information. Originally described as Gonyleptidae, Kury (2003) placed it as incertae sedis within Gonyleptoidea. Based on our study of a photograph of the holotype, we propose re-classifying it under Nomoclastidae, given the pedipalp morphology, absence of ornamentation in area I, and the shape of the dorsal shield type Zeta.

#### ***Quindina bella* Roewer, 1915**

*Quindina bella* Roewer, 1915: 128, fig. 14. — Roewer 1923: 564, fig. 707. — Soares and Soares 1948: 615. — Flórez and Sánchez-C 1995: 369. — Kury 2003: 97. — Pinto-da-Rocha and Bragagnolo 2017: 114, figs 5D, 8H, 14A, B.

**Distribution.** Tolima. Magdalena Valley montane forests (NT0136).

#### ***Quindina discolor* Pinzón & Pinto-da-Rocha, 2020**

*Quindina discolor* Pinzón & Pinto-da-Rocha, 2020: 542, figs 3, 4.

**Distribution.** Magdalena. Santa Marta montane forests (NT0159).

#### ***Quindina hermesi* Pinzón & Pinto-da-Rocha, 2020**

*Quindina hermesi* Pinzón & Pinto-da-Rocha, 2020: 537, figs 1, 2.

**Distribution.** Atlántico, Bolívar. Magdalena-Urabá moist forests (NT0137) and Guajira-Barranquilla xeric scrub (NT 1308).

#### ***Quindina marginata* (Roewer, 1963)**

*Deriacrus marginatus* Roewer, 1963: 68, fig. 42. — Flórez and Sánchez-C 1995: 368.

*Quindina marginata* — Pinto-da-Rocha and Bragagnolo 2017: 120, figs 7A, 9E, 14G, H.

**Distribution.** Huila. Magdalena Valley montane forests (NT0136).

#### **Stygnidae Simon, 1879**

##### ***Eutimesius ehippiatus* (Roewer, 1915)**

**Distribution.** Quindío. Magdalena Valley montane forests (NT0136).

***Eutimesius ornatus* (Roewer, 1943)**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Eutimesius simoni* Roewer, 1913**

**Distribution.** Putumayo. Napo moist forests (NT0142).

***Fortia jedi* Villarreal, Kury & Colmenares, 2022**

*Fortia jedi* Villarreal et al., 2022: 5, figs 1–11, 25–27.

**Distribution.** Magdalena. Santa Marta montane forests (NT0159).

***Fortia sith* Villarreal, Kury & Colmenares, 2022**

*Fortia sith* Villarreal et al., 2022: 8, figs 12–24, 28–30.

**Distribution.** Magdalena. Santa Marta montane forests (NT0159).

***Jabbastygnus huttoni* Kury & Villarreal, 2015**

*Jabbastygnus huttoni* Kury & Villarreal, 2015: 32, figs 4D–F, 17–19.

**Distribution.** Boyacá. Magdalena Valley montane forests (NT0136).

***Metaphareus albimanus* Roewer, 1912**

**Distribution.** Tolima. Magdalena Valley montane forests (NT0136).

***Niceforoielus assimilis* Mello-Leitão, 1941**

**Distribution.** Norte de Santander. Catatumbo moist forests (NT0108).

***Obidosus carnaval* (Villarreal-Manzanilla & Pinto-da-Rocha, 2006)**

*Protimesius carnaval* Villarreal-Manzanilla & Pinto-da-Rocha, 2006: 229, figs. 29–35, 44, 45.

*Obidosus carnaval* – Villarreal et al. 2019: 229, figs 29–35, 44, 45.

**Distribution.** Amazonas. Solimões-Japurá moist forests (NT0163).



**Remarks.** The type locality for this species is Moa River in the state of Acre, in Brazil. It was recorded for Colombia by Villarreal-M. and Pinto-da-Rocha (2006).

***Phareus antrophilus* Villarreal & Rodríguez, 2006**

*Phareus antrophilus* Villarreal & Rodríguez, 2006: 103, figs 1–9.

**Distribution.** Santander. Magdalena Valley montane forests (NT0136).

***Phareus raptator* (Gervais, 1844)**

*Goniosoma raptator* Gervais, 1844: 107 (vol. 3), pl. 47, figs 1, 1a.

*Liophareus mamillatus* Mello-Leitão, 1940: 309, fig. 10. — B. Soares 1945: 386.

*Colomphareus rugosus* Goodnight & Goodnight, 1943: 10, figs 32–35. — Flórez and Sánchez-C 1995: 368.

*Allophareus robustus* Roewer, 1963: 60, figs 26–28. — Flórez and Sánchez-C 1995: 368.

*Phareus raptator* — Simon 1879: 219. — Roewer 1912: 149. — 1913: 155, figs 68, 69. — 1923: 457, figs 475, 476. — Flórez and Sánchez-C 1995: 369. — Pinto-da-Rocha 1997: 220, figs 160–167, 489, 490, 582, 599. — Villarreal and Rodríguez 2006: 104.

**Distribution.** Cundinamarca. Cordillera Oriental montane forests (NT0118), Magdalena Valley montane forests (NT0136) and Northern Andean páramo (NT1006).

***Stygnus gertschi* (Roewer, 1963)**

**Distribution.** Cauca. Patía Valley dry forests (NT0225).

***Stygnus pectinipes* (Roewer, 1943)**

**Distribution.** Putumayo. Napo moist forests (NT0142).

***Stygnus simplex* (Roewer, 1913)**

**Distribution.** Putumayo. Napo moist forests (NT0142).

***Timesius vesicularis* (Gervais, 1844)**

**Distribution.** Department unknown.

**Samooidea Sørensen, 1886**  
**Stygnommatidae Roewer, 1923**

***Stygnomma fuhrmanni* Roewer, 1912**

**Distribution.** Antioquia. Cauca Valley montane forests (NT0109).

**Zalmoxoidea Sørensen, 1886**  
**Incertae sedis**

*Hevelia crucis* Kury, García & Ahumada-C., 2023  
*Hevelia crucis* Kury et al. 2023: 39, figs 1–4.

**Distribution.** Bolívar. Magdalena-Urabá moist forests (NT0137).

**Fissiphalliidae Martens, 1988**

***Fissiphallius spinulatus* Martens, 1988**

**Distribution.** Cundinamarca. Northern Andean paramo (NT1006).

***Fissiphallius sturmi* Martens, 1988**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

***Fissiphallius sympatricus* Martens, 1988**

**Distribution.** Cundinamarca. Magdalena Valley montane forests (NT0136).

**Icaleptidae Kury & Pérez, 2002**

***Icaleptes malkini* Kury & Pérez, 2002**

**Distribution.** Cesar. Santa Marta montane forests (NT0159).

**Kimulidae Pérez-González, Kury & Alonso-Zarazaga, 2007**

***Usatama infumatus* Kury, García & Medrano, 2019**

*Usatama* Kury, García, and Medrano 2019: 239, figs 1–4.

**Distribution.** Cundinamarca. Cordillera Oriental montane forests (NT0118).



### **Zalmoxidae Sørensen, 1886**

#### ***Minuides oedipus* Roewer, 1963**

*Minuides oedipus* Roewer, 1963: 48, figs 6–8; — Šilhavý 1978: 62. — Flórez and Sánchez-C 1995: 369.

**Distribution.** Huila. Magdalena Valley dry forests (NT0221).

**Remarks.** *Minuides oedipus* Roewer, 1963 treated as a Kimulidae in Kury's catalogue (Kury 2003); however, Pérez-González and Kury (2007) transferred *Minuides* Sørensen, 1932 to the Zalmoxidae.

#### ***Stygnoleptes analis* Banks, 1913**

**Distribution.** Department unknown.

#### ***Stygnoleptes crassus* (Sørensen, 1932)**

**Distribution.** Department unknown.

#### ***Timoleon crassipes* Sørensen, 1932**

**Distribution.** Department unknown.

### **Rationale for exclusions**

Two species, *Cosmetus flavopictus* Simon, 1880 and *Stygnoplus forcipatus* have traditionally been associated with the list of Colombian species, which we do not consider in our list. *Cosmetus flavopictus* was described for a locality in Darien, extrapolated by Kury (2003) as Colombian territory. Here, we follow the interpretation given by Medrano et al. (2021b) and consider not citing it as part of the Colombian species. *Stygnoplus forcipatus* Koch, 1845 was originally described without any locality data other than "Columb." (Koch 1845). Subsequently, it was recorded from Venezuela and its presence in the Colombian fauna was called into question (Villarreal-M. and Rodríguez 2004). Other species of the genus *Stygnoplus* are known from the Caribbean region, in the Venezuelan Cordillera de La Costa and the Lesser Antilles, with the exception of *S. longipalpus* (Goodnight & Goodnight, 1942), which occurs in the Guiana Shield. Recently, the only Amazonian species originally described in this genus by Pinto-da-Rocha and Tourinho (2012) were transferred to *Yapacana* Pinto-da-Rocha, 1997 (Villarreal et al. 2021b). We have decided not to include *S. forcipatus* in the list at this time due to lack of evidence of its presence in the country.





**Figure 13.** Selected Colombian harvestmen species, live specimens **A** *Barinas piragua* (Agoristenidae) from Bolívar **B** *Eucynorta quadripustulata* (Cosmetidae) from Bolívar **C** *Phalangodus cottus* (Cranidae) from Cundinamarca **D** *Phareicranaus* sp. (Cranidae) from Chocó **E** *Ampycus telifer* (Ampycidae) from Amazonas **F** *Quindina hermes* (Nomo-clastidae) from Bolívar **G** *Stygnomma* sp. (Stygnommatidae) from Tolima. Photographs by: Hugo Vides (**A**, **B**, **F**); Andrés García (**C**); Sarah Crews (**D**); Arthur Anker (**E**); Julio González / Entomopixel (**G**).

### Diversity

The Colombian opilionofauna of the suborder Laniatores is composed of 173 species, three of which are described in the present work. The 12 families listed in Colombia demonstrate the country's high taxonomic diversity for this group. The most diverse family is Cosmetidae with 67 species (Fig. 13B), followed by Cranidae with 40 species (Figs 13C, D, 14D). It is worth noting that a significant



number of Laniatores species in Colombia are poorly documented, with only vague records for the country or department, and their precise type localities remain unknown. Specifically, there are 16 species of Cosmetidae, five species of Cranidae, three species of Zalmoxidae, one species of Manosbiidae, two species of Nomoclastidae, one species *Incertae sedis*, and one species of Stygnidae with uncertain recorded information. These findings highlight our limited understanding of the diversity and distribution of this group in Colombia.

Most records come from a few departments, indicating a sampling bias. Approximately 22% of the species are known from the department of Cundinamarca (38 spp.) near the capital of the country, followed by Tolima and Boyacá with 13 species each. In contrast, all six departments of the Amazon


**Table 2.** Number of Opiliones species by family recorded in WWF terrestrial ecoregions from Colombia. When the locality is unknown or not precise enough, the abbreviation DNA is used, and the record is not assigned to any ecoregion. DNA = does not apply.

WWF Ecoregion	Code (WWF)	Families	Species
Magdalena Valley montane forests	NT0136	Agoristenidae	6
		Cosmetidae	25
		Cranidae	12
		Fissiphalliidae	2
		Manosbiidae	5
		Nomoclastidae	4
		Stygnidae	6
Cordillera Oriental montane forests	NT0118	Agoristenidae	4
		Cosmetidae	10
		Cranidae	1
		Kimulidae	1
		Stygnidae	1
Northern Andean páramo	NT1006	Cosmetidae	10
		Cranidae	1
		Fissiphalliidae	1
		Stygnidae	1
Cauca Valley montane forests	NT0109	Cranidae	9
		Manosbiidae	1
		Stygnomatidae	1
Guajira-Barranquilla xeric scrub	NT1308	Agoristenidae	3
		Cosmetidae	4
		Cranidae	3
		Nomoclastidae	1
Chocó-Darién moist forests	NT0115	Cosmetidae	7
		Cranidae	2
		Ampycidae	1
Napo moist forests	NT0142	Agoristenidae	1
		Cosmetidae	2
		Cranidae	1
		Stygnidae	3

WWF Ecoregion	Code (WWF)	Families	Species
Solimões-Japurá moist forests	NT0163	Cosmetidae	3
		Ampycidae	2
		Nomoclastidae	1
		Stygnidae	1
Santa Marta montane forests	NT0159	Agoristenidae	1
		Icaleptidae	1
		Nomoclastidae	1
		Stygnidae	2
Magdalena Valley dry forests	NT0221	Agoristenidae	1
		Cosmetidae	1
		Cranidae	2
		Zalmoxidae	1
Northwestern Andean montane forests	NT0145	Cosmetidae	1
		Cranidae	2
Apure-Villavicencio dry forests	NT0201	Cranidae	2
		Ampycidae	1
Patía Valley dry forests	NT0225	Cranidae	1
		Manaosbiidae	1
		Stygnidae	1
Catatumbo moist forests	NT0108	Agoristenidae	1
		Manaosbiidae	1
		Stygnidae	1
Magdalena-Urabá moist forests	NT0137	Agoristenidae	1
		Cosmetidae	1
		Incertae sedis	1
		Nomoclastidae	1
Sinú Valley dry forests	NT0229	Cosmetidae	2
Amazon-Orinoco-Southern Caribbean mangroves	NT1401	Agoristenidae	1
		Cosmetidae	1
Caqueta moist forests	NT0107	Cosmetidae	1
		Ampycidae	1
		Nomoclastidae	1
Purus várzea	NT0156	Cosmetidae	2
Eastern Cordillera real montane forests	NT0121	Cranidae	1
Iquitos várzea	NT0128	Cosmetidae	1
Llanos	NT0709	Cosmetidae	1
Unknown Locality	DNA	Cosmetidae	21
		Cranidae	6
		Manaosbiidae	1
		Nomoclastidae	1
		Stygnidae	1
		Zalmoxidae	3

region have a combined total of only 15 recorded species. This exemplifies the limited knowledge we have on the diversity of harvestmen in the Colombian Amazon region. Of the 173 species of Laniatores known in Colombia, 35 are recorded without a specific locality; 28 of which have no information other than their presence in Colombia, and only seven have been located at a spe-






**Figure 14.** Laniatores species described from Colombia **A** the pie chart shows authors who described two or more species. For each author, every described species was counted, regardless of whether the authorship of the species is shared with other authors. For example, *Usatama infumatus* was described by Kury, García & Medrano, 2019, and was counted as a species for each author. Authors with one species are included in the others category **B** accumulation curve of Laniatores species described from Colombia. Pink: descriptions made between 1833 and 1911; ochre: 1912–1963; green: 1964–2023 **C** Laniatores species in RUNAP **D** families of Laniatores species described from Colombia.

cific department (Table 2). The remaining departments have eight or fewer recorded species each, and eight departments have no records (Arauca, Caldas, Guainía, Guaviare, Risaralda, San Andrés y Providencia, Sucre, and Vichada).

Olson et al. (2001) recognized 35 ecoregions in Colombia, but only 22 of them have Laniatores species recorded (Table 2). The ecoregion with the highest number of species is the Magdalena Valley montane forests, with a total of 59 species. The ecoregions with the fewest species are Eastern Cordillera real montane forests, Iquitos várzea, Llanos, and Purus várzea, each with one species (Figs 15, 16). Only 21 species are found in more than one ecoregion, with *Eucynorta quadripustulata* (Fig. 13B) being the most widely recorded species, occurring in five ecoregions.




**Figure 15.** Geographical distribution of the families **A** Cosmetidae and **B** Cranidae in Colombia. Colored polygons follow the regionalization of the Neotropical region proposed by WWF (Olson et al. 2001).

The Colombian NSPA lists 1,486 protected zones, of which 60 are NNP and FFS. The remaining areas are designated with some other category of protection (RUNAP 2022). Regarding Opiliones Laniatores, 36 species have been recorded from 29 areas, totaling 47 records, and 11 species have been recorded in seven NNP or FFS. The protection category with the highest number of the recorded species is RDIM, with 16 species, followed by NPFR with ten species, NNP with eight species, FFS with four species, PNR with three species, RFPR with two species, and RNSC with one species.

Similar to other neotropical countries, the taxonomic knowledge of harvestmen in Colombia is primarily based on the contributions of foreign authors. Remarkably Carl F. Roewer described almost half of the species known of the country. Therefore, for a pragmatic approach we divided the history into three periods: the first period prior to Roewer's contributions, the second period in which most of the descriptions took place, led by Roewer's works but with contributions from other authors, and finally, the third, post-Roewerian or modern period, with contributions from a diverse group of authors, including local authors for the first time. Our goal is not to present a detailed chronology, but rather to provide a general overview of the historical panorama and the individual contribution of the authors to the knowledge of the Colombian fauna. For a detailed chronicle and an in-depth look at the early history of opilionology, we recommend referring to Kury (2010), who presented an excellent compilation and analysis of papers between 1758 and 1804. Figure. 14B shows a timeline






**Figure 16.** Geographical distribution of **A** Gonyleptoidea (Ampycidae, Manaosbiidae, Nomoclastidae and Stygnidae) and **B** Samooidea (Kimulidae, Stygnommatidae) and Zalmoxoidea (Icaleptidae, Fissiphalliidae, Zalmoxidae) in Colombia. Colored polygons follow the regionalization of the Neotropical region proposed by WWF (Olson et al. 2001).

of descriptions of armored harvestmen species in Colombia and the contributions by author and periods.

Perty (1833) described *Sibambea cincta* (Perty, 1833), as the first armored harvestmen species known for Colombia. This is the starting point for the ancient or Pre-Roewerian period (1833–1880), during which 19 species were described by four authors, highlighting the contribution by French authors Eugène Simon (nine species described) and Paul Gervais (seven described species). The second period or so-called Roewerian period (1912–1963), is characterized by a substantial increase in the number of species described, thanks to the prolific work of the German author Roewer (71 Colombian species described between 1912 and 1963). Other important authors from this period include Mello-Leitão, Sørensen and Clarence J. and Marie L. Goodnight, totaling 27 described species. After Roewer's death, there was a period of taxonomic inactivity for the Colombian fauna, until Martens (1988) published a paper on Colombian harvestmen, marking the beginning of the Modern or Post-Roewerian period (1988–present). Since then, contributions have become more frequent, particularly from Latin American authors, such as Brazilian authors A. B. Kury (22 spp.) and R. Pinto-da-Rocha (10 spp.) and Venezuelan author O. Villarreal (18 spp.), as well as local authors, highlighting A. F. García (17 spp.) and M. Medrano (11 spp.) (Fig. 14A).





**Figure 17.** Geographical distribution of the families **A** Cosmetidae and **B** Cranidae in Colombia. Colored polygons represent the RUNAP areas. Civil Society Nature Reserve (CSNR), Flora and Fauna Sanctuary (FFS), National Protective Forest Reserve (NPFR), Regional Natural Park (RNP), Regional Protective Forest Reserve (RPFR), National Natural Parks (NNP), Regional Integrated Management Districts (RDIM).

## Discussion


### Diversity of Colombian Opiliones

Colombia is recognized as a megadiverse country, renowned for its numerous protected natural areas compared to other neotropical countries. Despite ongoing conservation efforts, there is a need to increase the knowledge on biodiversity in these areas to support their proper management and handling. Unhappily, the information available on Laniatores in these areas is still limited, despite the growing number of arachnologists in the region and the country, and the increasing research focus on local fauna (Fig. 14C).

The central and eastern cordillera are the most populated and accessible areas of Colombia. They are home to the largest universities and biological collections in the country and, consequently, the region with the highest number of descriptions and records of Laniatores (Figs 15, 16). Local researchers have made significant contributions to the study of Laniatores in the last decade, including the review of some genera such as *Rhaucus* or *Eulibitia* (García and Kury 2017; Medrano and Kury 2017) and description of some new species (e.g., Kury et al. 2019; García and Ahumada-C. 2022).

In contrast, the peripheral zones, like the Amazon, the Pacific, or the Caribbean, have received less attention from researchers. Nevertheless, recent efforts





**Figure 18.** Geographical distribution of **A** the superfamilies Gonyleptoidea (Agoristenidae, Ampycidae, Manaosbiidae, Nomoclastidae and Stygnidae) and **B** Samooidea (Kimulidae, Stygnommatidae) and Zalmoxoidea (Icaleptidae, Fissiphalliidae, Zalmoxidae) in Colombia. Colored polygons represent the RUNAP areas. Flora and Fauna Sanctuary (FFS), National Protective Forest Reserve (NPFR), National Natural Parks (NNP), Regional Integrated Management Districts (RDIM).

have been made to strengthen research in these regions. For instance, a school of arachnology at the Universidad del Valle, has emerged to study the Laniatores of Valle del Cauca in the Pacific region. Similarly, in the Caribbean, harvestmen have been the focus of some degree work, and recent collections have yielded specimens that have been added to the country's biological collections, resulting in the descriptions of new species and new records (e.g., Pinzón and Pinto-da-Rocha 2020; Ahumada-C. et al. 2022; Kury et al. 2023).

Therefore, we believe that although the national inventory of Laniatores is far from being complete, further work on the local fauna will increase our knowledge of the group over time. Regarding the Colombian protected areas, the RNSC is the area with the lowest number of records of Laniatores (one species), while RDIM presents the highest number of records (16 species), probably because it is the only protected area in the country conceived as a rational use mode (Molina 2013), with research permits that are easier to access. However, when considering the total number of records for Laniatores across all categories of protected areas, studies are also scarce (Figs 17, 18). The implementation of strategies that facilitate the necessary regulatory procedures to collect specimens in protected natural areas, as well as increased investment in basic science projects, could help to generate a greater understanding of Colombian biodiversity in these areas. A successful case is the study by García

and Medrano (2015) in the Reserva Natural Rio Ñambí, the only faunal study on harvestmen conducted in a protected natural area, resulting in the discovery of approximately 28 species new to science, demonstrating the hidden diversity in these areas, and the importance of inventories or taxonomic studies on the protected natural areas.

Continued exploration of ecosystems is imperative to search for Opiliones and achieve reliable taxonomic identifications, which are essential to determine the true distribution of the species. Such data is important in establishing their threat category, according with the criteria from the International Union for Conservation of Nature (IUCN). This should be a goal for arachnology, given the high endemism exhibited by these organisms and the significant anthropogenic pressures to which they are exposed.

## Taxonomic accounts

*Phalangodus* is an Andean cranaid genus whose species could be considered examples of short-range endemism, as defined by Harvey (2002). The distribution of *Phalangodus andresi* sp. nov. is within the area of generic distribution and represents the first case of sympatry in the genus, as it was collected in close proximity to *P. anacosmetus*, a widely distributed species found in the highlands of the department of Cundinamarca (Villarreal and García 2016).

*Phalangodus andresi* sp. nov. possesses all generic diagnostic characters presented by Villarreal and García (2016), and can be diagnosed by its smaller size, by having dorsal scutum tuberculated (shared only with *P. palpiconus*), being distinguished by the type of ornamentation of the femur IV of the males (refer to the diagnosis for further details). Unfortunately, of the type locality for *P. palpiconus* was erroneously attributed to Chile, which is unlikely due to the distribution pattern of the genus and even the family, as discussed by Hara et al. (2014), making impossible to establish any biogeographic relationship between both species.

Moreover, the family Ampycidae is known to have a little-explored diversity, primarily in Ecuador and Colombia. However, the taxonomic assignment of new species is a challenge for taxonomists due to the paucity of old descriptions and poor generic diagnosis. In addition, many species exhibit similar external morphology, and for more species, genitalia remain unknown, with just some exceptions (e.g., Villarreal and Kury 2012; Kury and Quintero 2014; Tourinho and Mendes 2014; Hara et al. 2017). Despite these challenges, recent studies on some species have shed light on the external morphology and genitalia of the subfamily. Further research is necessary to address the taxonomic uncertainties and increase our knowledge of the diversity of Ampycidae.

## Acknowledgments

To Eduardo Flórez, Daniela Martínez and Sebastian Galvis for their collaboration during the visit by DAC and OV to the arachnological collection of ICN. To Adriano B. Kury, for all the facilities in the AracnoLab (MNRJ) work during the visit by DAC, for maintaining the OmniPaper Project and WCO Lite, and for his excellent role as Subject Editor for this article and for allowing us to access photographs of the type specimens of *Ampycella* and *Paraphalangodus synacanthus* as well



as the constant feedback. To Marco R. Hara for the feedback regarding the identity of *Phalangodus andresi* sp. nov. in relation to *P. palpiconus*. We would like to express our gratitude to the Vice-Rector's Office for Research of the Universidad del Quindío for their invaluable assistance in reviewing the English writing of our manuscript. To Andrés F. García for the Scanning Electron Microscopy (SEM) photographs used in Fig. 7. The SEM photographs were taken at the Microscopy Platform Rudolf Barth/IOC-Fiocruz. OV thanks Pío Colmenares (AMNH) for coordinating the donation of optical equipment and supplies to MIZA through the Cooperation Agreement between MIZA-AMNH. The following photographers kindly provided the use of their photographs of live harvestmen for Fig. 13: Andrés García, Arthur Anker, Hugo Vides, Julio González /Entomopixel, and Sarah Crews. To Miguel Medrano, Ligia Benavides, and one anonymous referee, thank you for valuable suggestions and comments that greatly improved the present work; to Nathalie Yonow for her editorial revision work. To Rodrigo Baos for his kindness in San Pedro, Buenaventura, Valle del Cauca.

## Additional information

### Conflict of interest

The authors have declared that no competing interests exist.

### Ethical statement

No ethical statement was reported.

### Funding

This study has been partly supported by the Vice-Rector's Office for Research of the Universidad del Quindío for LDS through the founding of the publication costs. Idea Wild funded photographic equipment for OV in the framework of the project ID. VILLVENE0122. Universidad de Cartagena funded the DAC's visit to MNRJ (Resolution 01787 of 2020).

### Author contributions

OVM conceived the project, structured an initial draft of the manuscript, identified the new species, and contributed to their description and illustration, as well as with the list of species. DAC participated in the identification of the species, elaboration of the descriptions and illustrations, worked on the list of species, and collaborated significantly in the final writing of the manuscript. LDS collected two of the new species, participated in the identification of the species, contributed to the imaging, coordinated the making of the maps, and contributed to the overall writing of the manuscript. All authors contributed equally by writing results and discussion and agreed with the final version of the manuscript.

### Author ORCIDs

Oswaldo Villarreal  <https://orcid.org/0000-0001-5355-3723>

Daniela Ahumada-C.  <https://orcid.org/0000-0002-4182-5143>

Leonardo Delgado-Santa  <https://orcid.org/0000-0002-9714-6924>

### Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

## References

- Acosta LE, Pérez-González A, Tourinho A (2007) Methods for taxonomic study. In: Pinto-da-Rocha R, Machado G, Giribet G (Eds) *Harvestmen: The Biology of Opiliones*. Harvard University Press, Cambridge and London, England, 494–510.
- Adis J, Bonaldo AB, Brescovit AD, Bernati R, Cokendolpher JC, Condé B, Kury AB, Lourenço WR, Mahnert V, Pinto-da-Rocha R, Platnick NI, Reddell JR, Rheims CA, Rocha LS, Rowland JM, Weygoldt P, Woas S (2002) Arachnida at “Reserve Ducke”, Central Amazonia/Brazil. *Amazoniana* 17: 1–14.
- Ahumada-C. D, García AF, Navas-S GR (2020) The spiny agoristenid genus *Barinas* (Arachnida: Opiliones), with the description of a new species from the Colombian Caribbean. *Arachnology* 18(6):632–641. <https://doi.org/10.13156/arac.2020.18.6.632>
- Ahumada-C. D, Medrano M, Segovia-Paccini A (2022) Redescription and geographic distribution of the type species of *Eucynorta* (Arachnida, Opiliones, Cosmetidae). *Studies on Neotropical Fauna and Environment* 1–10. <https://doi.org/10.1080/01650521.2022.2076964>
- Andrade-C MG (2011) Estado del conocimiento de la Biodiversidad en Colombia y sus amenazas. Consideraciones para fortalecer la interacción Ciencia-Política. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 35(137): 491–507.
- Barriga JC, Martínez-Torres D, López-Orozco CM, Villarreal O, Murcia M (2019) Artrópodos terrestres de las cuevas y cavernas de El Peñón (Andes), Santander, Colombia. In: Lasso CA, Barriga JC, Fernández-Auderset, Y (Eds) *Biodiversidad subterránea y epigea de los sistemas cársticos de El Peñón (Andes), Santander, Colombia*. VII. Serie Fauna Silvestre Neotropical, Instituto de Investigación de Recursos Bioológicos Alexander von Humboldt, Bogotá, D.C. Colombia, 99–154.
- Bragagnolo C, Pinto-da-Rocha R (2003) Diversidade de Opiliões do Parque Nacional da Serra Dos Órgãos, Rio de Janeiro, Brasil (Arachnida: Opiliones). *Biota Neotropica* 3(1): 1–20. <https://doi.org/10.1590/S1676-06032003000100009>
- Centore P (2016) sRGB Centroids for the ISCC-NBS Colour System. Self Published, 21 pp.
- CONPES [Consejo Nacional de Política Económica y Social República de Colombia] (2021) Departamento Nacional de Planeación, Bogotá D.C., Colombia, 114 pp.
- DaSilva MB, Gnaspini PA (2010) Systematic revision of Goniosomatinae (Arachnida: Opiliones: Gonyleptidae), with a cladistic analysis and biogeographical notes. *Invertebrate Systematics* 23(6): 530–624. <https://doi.org/10.1071/IS09022>
- De Moya N, Martínez NJ, García AF (2021) Diversidad taxonómica de opiliones en la vertiente occidental de la Sierra Nevada de Santa Marta, Magdalena. Colombia. *Boletín Científico Museo de Historia Natural Universidad de Caldas* 25: 157–180. <https://doi.org/10.17151/bccm.2021.25.2.10>
- ESRI (2022) Environmental Systems Research Institute – ESRI. ArcGIS desktop. [Available online:] <https://www.esri.com> [Accessed 03 July 2022]
- Flórez E, Sánchez-C H (1995) La diversidad de los arácnidos de Colombia, aproximación inicial. In: Rangel IO (Ed.) *Colombia diversidad biótica*. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, IMANI: Bogotá D.C., Colombia, 327–345.
- García AF (2014) Primeros registros de Ampycinae Kury, 2003 (Opiliones, Gonyleptidae) en Colombia. *Revista Iberica de Aracnologia* 25: 93–95.
- García AF, Ahumada-C. D (2018) Completing the puzzle: Another species of *Rhaucus* Simon, 1879 (Arachnida: Opiliones: Cosmetidae) from Colombia. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales* 42(163): 200–206. <https://doi.org/10.18257/raccefyn.611>



- García AF, Ahumada-C. D (2022) Taxonomic notes on *Barinas*: A new generic synonym, a new cave-dwelling species, and new records from Colombia (Arachnida, Opiliones, Agoristenidae). *Evolutionary Systematics* 6(1): 1–7. <https://doi.org/10.3897/evol-syst.6.78123>
- García AF, Kury AB (2017) Taxonomic revision of the Andean harvestman genus *Rhaucus* Simon, 1879 (Arachnida, Opiliones, Cosmetidae). *Zootaxa* 4338(3): 401–404. <https://doi.org/10.11646/zootaxa.4338.3.1>
- García AF, Kury AB (2020) The Neotropical harvestman genus *Vima* Hirst, with description of a new species from Colombia (Arachnida: Opiliones: Agoristenidae). *The Journal of Arachnology* 48(1): 67–76. <https://doi.org/10.1636/0161-8202-48.1.67>
- García AF, Medrano M (2015) Orden Opiliones. In: Flórez E, Romero C, López DS (Eds) *Los Artrópodos de la reserva Natural Río Ñambí*. Instituto de Ciencias Naturales, Universidad Nacional de Colombia: Bogotá D.C., Colombia, 109–129.
- García AF, Pastrana-M MR (2021) A remarkable new species of Agoristenidae (Arachnida, Opiliones) from Córdoba, Colombia. *Papéis Avulsos de Zoologia* 61: e20216127. [8 pp] <https://doi.org/10.11606/1807-0205/2021.61.27>
- García AF, Villarreal O (2020) Description of a new species of *Leptostygnus* Mello-Leitão, 1940 and notes on the male genitalia in the subfamily Leiosteninae (Opiliones: Agoristenidae). *Studies on Neotropical Fauna and Environment* 55(1): 1–15. <https://doi.org/10.1080/01650521.2020.1724496>
- García AF, González A, Gutiérrez M (2022) New records and a new cave-dwelling species of Agoristenidae (Arachnida, Opiliones) from Colombia. *Zoosystematics and Evolution* 98(1): 55–63. <https://doi.org/10.3897/zse.98.78202>
- Guerrero EL (2019) Los opiliones (Arachnida: Opiliones) de las áreas protegidas de la provincia de Buenos Aires y la Ciudad Autónoma de Buenos Aires, Argentina. *Revista de la Sociedad Entomológica Argentina* 78(4): 4–13. <https://doi.org/10.25085/rsea.780402>
- Hadley A (2015) CombineZP. Image stacking software. 2015. <http://www.hadleyweb.pwp.blueyonder.co.uk/CZP/Installation.htm> [Accessed 03 July 2022]
- Hara MR, Pinto-da-Rocha R, Villarreal O (2014) Revision of the cranaid genera *Phalangodus*, *Iquitosa* and *Aguaytiella* (Opiliones: Laniatores: Gonyleptoidea). *Zootaxa* 3814(4): 567–580. <https://doi.org/10.11646/zootaxa.3814.4.8>
- Hara MR, Bragagnolo C, Pinto-da-Rocha R (2017) Revision of *Thaumatochranaus* Roewer, 1932, with description of two new species (Opiliones: Laniatores: Gonyleptoidea). *Zootaxa* 4254(4): 457–472. <https://doi.org/10.11646/zootaxa.4254.4.3>
- Harrington B (2004–2005) Inkscape. [Available online:] <http://www.inkscape.org/> [Accessed 05 December 2022]
- Harvey MS (2002) Short-range endemism amongst the Australian fauna: Some examples from non-marine environments. *Invertebrate Systematics* 16(4): 555–570. <https://doi.org/10.1071/IS02009>
- Koch CL (1845) *Die Arachniden getreu nach der Natur abgebildet und beschrieben*. Vol. 7: 1–130, 36 pls, CH Zeh, Nuremberg, 166 pp.
- Kury AB (2003) Annotated catalogue of the Laniatores of the New World (Arachnida, Opiliones). *Revista Iberica de Aracnologia* 1: 1–337.
- Kury AB (2010) Opilionological Record – a chronicle of harvestman taxonomy. Part 1: 1758–1804. *The Journal of Arachnology* 38(3): 521–529. <https://doi.org/10.1636/Ha10-23.1>
- Kury AB, Barros C (2014) A new genus and eight new species of Amazonian cosmetines (Opiliones, Laniatores, Cosmetidae). *Zoological Studies (Taipei, Taiwan)* 53(1): 1–46. <https://doi.org/10.1186/s40555-014-0024-4>



- Kury AB, Medrano M (2016) Review of terminology for the outline of dorsal scutum in Laniatores (Arachnida, Opiliones). *Zootaxa* 4097(1): 130–134. <https://doi.org/10.11646/zootaxa.4097.1.9>
- Kury AB, Quintero D (2014) Characterization of the genus *Hernandarioides* FO Pickard-Cambridge, 1905 (Opiliones, Gonyleptidae, Ampycinae). *Zootaxa* 3838(2): 242–246. <https://doi.org/10.11646/zootaxa.3838.2.9>
- Kury AB, Villarreal MO (2015) The prickly blade mapped: Establishing homologies and a chaetotaxy for macrosetae of penis ventral plate in Gonyleptoidea (Arachnida, Opiliones, Laniatores). *Zoological Journal of the Linnean Society* 174(1): 1–46. <https://doi.org/10.1111/zoj.12225>
- Kury AB, García AF, Medrano M (2019) A new genus of Kimulidae; first record of the family from Colombia (Opiliones, Grassatores). *Comptes Rendus Biologies* 342(5–6): 236–244. <https://doi.org/10.1016/j.crv.2019.07.003>
- Kury AB, Mendes AC, Cardoso L, Kury MS, de Andrade A, Giribet G, Cruz-López JA, Longhorn SJ, Medrano M, Kury IS, Kury M (2022) World Catalogue of Opiliones. WCO-Lite version 2.4.0. 2022. [Available online:] <https://wcolite.com/> [Accessed 04 December 2022]
- Kury AB, García AF, Ahumada-C. D (2023) A new genus of Zalmoxoidea from Colombia (Arachnida: Opiliones: Grassatores). *Journal of Arachnology* 51: 37–45. <https://doi.org/10.1636/JoA-S-21-070>
- Martens J (1988) Fissiphalliidae, a new family of South American laniatorean harvestmen (Arachnida: Opiliones). *Journal of Zoological Systematics and Evolutionary Research* 26(2): 114–127. <https://doi.org/10.1111/j.1439-0469.1988.tb00303.x>
- Medrano MA, Kury AB (2017) Taxonomic revision of the Andean genus *Eulibitia* Roewer, 1912 (Arachnida, Opiliones, Cosmetidae), with the description of five new species. *European Journal of Taxonomy* 357: 1–55. <https://doi.org/10.5852/ejt.2017.357>
- Medrano M, Ázara L, Kury AB (2019) Rediscovery of *Eulibitia ectroxantha* (Mello-Leitão, 1941) and synonymy of *Sphalerocynorta* Mello-Leitão, 1933 (Opiliones: Cosmetidae). *Comptes Rendus Biologies* 342(9–10): 345–350. <https://doi.org/10.1016/j.crv.2019.11.002>
- Medrano M, Ázara L, Kury AB (2020) The short-legged Andean cosmetids revisited: The genus *Libitia* Simon, 1879 with description of two new species (Opiliones, Cosmetidae). *European Journal of Taxonomy* 634(634): 1–25. <https://doi.org/10.5852/ejt.2020.634>
- Medrano M, Kury AB, Mendes MC (2021a) Morphology-based cladistics splinters the century-old dichotomy of the pied harvestmen (Arachnida: Gonyleptoidea: Cosmetidae). *Zoological Journal of the Linnean Society* 195(2): 585–672. <https://doi.org/10.1093/zoolinnean/zlab043>
- Medrano M, Kury AB, Martínez L (2021b) A fresh look at *Cosmetus* Perty, 1833 (Opiliones: Cosmetidae), with new synonymies and description of two new species from Colombia. *Zootaxa* 5004(3): 430–446. <https://doi.org/10.11646/zootaxa.5004.3.2>
- Medrano M, García AF, Kury AB (2022) *Rhaucoides* Roewer, 1912, an Andean genus with fused tarsomeres: revision with a new generic synonymy and two new species (Opiliones: Cosmetidae: Metergininae). *The Journal of Arachnology* 50(2): 191–218. <https://doi.org/10.1636/JoA-S-21-028>
- Molina E (2013) Distritos de manejo integrado: Estrategia de conservación y utilización sostenible de la biodiversidad. *Revista Ingeniería de los Recursos Naturales del Ambiente*. <http://revistaeidenar.univalle.edu.co/revista/ejemplares/12/d.htm> [Accessed 10 April 2023]
- Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GVN, Underwood EC, D’Amico JA, Strand HE, Morrison JC, Loucks CJ, Allnutt TF, Lamoreux JF, Ricketts



- TH, Itoua I, Wettengel WW, Kura Y, Hedao P, Kassem K (2001) Terrestrial Ecoregions of the World: A New Map of Life on Earth. *Bioscience* 51(11): 933–938. [https://doi.org/10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2)
- Orrico VGD, Kury AB (2009) A cladistic analysis of the Stygnicraninae Roewer, 1913 (Arachnida, Opiliones, Cranida) – Where do longipalp cranids belong? *Zoological Journal of the Linnean Society* 157(3): 470–494. <https://doi.org/10.1111/j.1096-3642.2009.00543.x>
- Perafán C, Sabogal A, Moreno-González JA, García-Rincón A, Luna-Sarmiento D, Romero-Ortiz C, Flórez E (2013) Diagnóstico del estado actual de la fauna de arácnidos y de su gestión en Colombia. In: Rueda-Ramírez D, Torrado-León E, Becerra EH (Comp) *Memorias Congreso Colombiano de Entomología, 40 Congreso Socolen, Sociedad Colombiana de Entomología – Socolen: Bogotá, D.C., Colombia*, 308–335.
- Pérez-González A, Kury AB (2007) Kimulidae Pérez González, Kury Alonso-Zarazaga In: Pinto-da-Rocha R, Machado G, Giribet G (Eds) *Harvestmen: the biology of the Opiliones*, Harvard University Press: Cambridge and London, England, 207–209.
- Perty JAM (1833) *Delectus animalium articulorum, quae in itinere per Brasiliam annis MDCCCXVII–MDCCCXX [1817–1820] jussu et auspiciis Maximiliani Josephi I Bavariae Regis augustissimi peracto, collegerunt Dr. J. B. de Spix et Dr. C. F. Ph. de Martius [1830–1834]*, Friedrich Fleischer, Monachii (= München), 1833. <https://doi.org/10.5962/bhl.title.102991>
- Pinto-da-Rocha R, Bonaldo AB (2011) Species relationships in the Neotropical genus *Phareicranaus* Roewer 1913 (Opiliones: Cranida): two new species and new data from Penial morphology. *Zootaxa* 3135(1): 1–34. <https://doi.org/10.11646/zootaxa.3135.1.1>
- Pinto-da-Rocha R, Bragagnolo C (2017) Cladistic analysis of the family Nomoclastidae with descriptions of a new genus and eight new species (Opiliones, Laniatores). *Invertebrate Systematics* 31(1): 91–123. <https://doi.org/10.1071/IS15050>
- Pinto-da-Rocha R, Tourinho AL (2012) Two new genera, ten new species and new records of Amazonian Stygnidae Simon, 1879 (Opiliones: Laniatores). *Zootaxa* 3340(1): 1–28. <https://doi.org/10.11646/zootaxa.3340.1.1>
- Pinzón MCA, Pinto-da-Rocha R (2020) Two new Colombian harvestmen of the genus *Quindina* Roewer, 1914 (Opiliones: Nomoclastidae). *Zootaxa* 4748(3): 531–547. <https://doi.org/10.11646/zootaxa.4748.3.7>
- Pinzón-M. C, Townsend VR, Martínez HN (2017) New record for the genus *Platymessa* Mello-Leitão, 1941 in Colombia, with the description of a new species (Opiliones, Cosmetidae). *ZooKeys* 665: 59–70. <https://doi.org/10.3897/zookeys.665.11371>
- Pinzón C, Damron B, Pinto-da-Rocha R (2021) Three new species of Cosmetidae C.L. Koch, 1839 from South America (Opiliones: Cosmetidae). *Zootaxa* 4984(1): 203–217. <https://doi.org/10.11646/zootaxa.4984.1.16>
- RUNAP (2022) Reporte de Áreas inscritas en el Registro Único Nacional de Áreas Protegidas–RUNAP.
- Soares BAM, Soares HEM (1970) Opiliões de Itatiaia (Opiliones: Gonyleptidae, Phalangodidae). *Revista Brasileira de Biologia* 30(3): 339–350.
- Tourinho AL, Mendes A (2014) New species of the harvestmen *Hutamaia* (Laniatores: Gonyleptidae: Ampycinae) and generic diagnosis. *Zoologia* 31(5): 463–474. <https://doi.org/10.1590/S1984-46702014000500006>
- Villarreal MO, García AF (2016) Unveiling the diversity of *Phalangodus* Gervais, 1842 (Opiliones: Cranida): descriptions of four new species from Colombia. *European Journal of Taxonomy* 242(242): 1–141. <https://doi.org/10.5852/ejt.2016.242>



- Villarreal MO, García AF (2021) On the phylogenetic relationships of *Muscopilio*, a new Andean genus of basibiont harvestmen (Opiliones: Agoristenidae). *Zoologischer Anzeiger* 292: 150–162. <https://doi.org/10.1016/j.jcz.2021.03.006>
- Villarreal MO, Kury AB (2012) *Licornus* Roewer, 1932: newly transferred to Ampycinae and first record of the family Gonyleptidae (Opiliones: Laniatores) from Venezuela. *Zootaxa* 3544(1): 71–78. <https://doi.org/10.11646/zootaxa.3544.1.6>
- Villarreal O, Rodríguez C (2006) Nuevos registros del género *Phareus* Simon, 1879 en Colombia, con la descripción de una nueva especie (Opiliones, Stygnidae). *Boletín de la SEA* 38: 103–105.
- Villarreal MO, Kury AB, Pinto-da-Rocha R (2015) The poorly known genus *Ventrifurca* Roewer, 1913 revisited (Opiliones: Cranidae). *Zoological Studies (Taipei, Taiwan)* 54(1): 1–18. <https://doi.org/10.1186/s40555-015-0117-8>
- Villarreal O, García AF, Kury AB (2021a) Fine-tuning the diversity in four families of Gonyleptoidea (Arachnida: Opiliones) in Venezuela. *Revista Mexicana de Biodiversidad* 92(0): 1–22. <https://doi.org/10.22201/ib.20078706e.2021.92.3633>
- Villarreal O, Kury AB, Colmenares PA (2021b) Contributions to the taxonomy of some Amazonian Stygnidae (Opiliones: Laniatores: Gonyleptoidea). *Zootaxa* 4984(1): 218–227. <https://doi.org/10.11646/zootaxa.4984.1.17>
- Villarreal O, Kury AB, Colmenares PA (2022) New genus with two new species of Colombian harvestmen (Opiliones: Stygnidae: Stygninae). *American Museum Novitates* 3991(3991): 1–14. <https://doi.org/10.1206/3991.1>
- Villarreal-M. O, Rodríguez CJ (2004) Descripción de una nueva especie y dos nuevos registros del género *Stygnoplus* (Opiliones, Stygnidae) para Venezuela. *Revista Iberica de Aracnología* 10: 179–184.
- Villarreal-Manzanilla O, Pinto-da-Rocha R (2006) Five new species of *Protimesius* from Brazil (Opiliones: Stygnidae). *Zootaxa* 1325(1): 219–233. <https://doi.org/10.11646/zootaxa.1325.1.14>

## Supplementary material 1

### List of Laniatores species for Colombia

Authors: Osvaldo Villarreal, Daniela Ahumada-C., Leonardo Delgado-Santa

Data type: xlsx

Explanation note: List of Laniatores species for Colombia, with their registration source, distribution associated with WWF biogeographic ecoregions and their presence in protected areas, National System of Protected Areas.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/zookeys.1175.102485.suppl1>